arXiv:1304.3862v1 [math-ph] 13 Apr 2013

ABSOLUTELY CONTINUOUS SPECTRUM FOR RANDOM SCHRÖDINGER OPERATORS ON THE FIBONACCI TREE-STRIP

CHRISTIAN SADEL

Abstract. The Fibonacci tree is a special non-regular tree. The tree-strip is the cross product with a finite graph. We prove that for small disorder the Anderson model on this tree-strip has purely absolutely continuous spectrum in a certain set.

1. Introduction

The Fibonacci trees are infinite trees associated to the substitution matrix $S = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 1 \end{smallmatrix}\right)$. This means that the trees are constructed starting from a root and going from generation to generation in the following way. Each vertex x of the tree has either the label $l(x) = 1$ or $l(x) = 2$, each vertex with label 1 has one child with label 1 and each vertex with label 2 has one child of label 1 and one child of label 2. So each vertex of label 1, except for possibly the root, has 2 neighbors (one parent and 1 child), and each vertex of label 2, except for possibly the root, has 3 neighbors (one parent and 2 children). The label of the root determines the tree up to tree isomorphism (see Figure 1).

Let $\mathbb{T}^{(1)}$ denote the Fibonacci tree with root label 1, and $\mathbb{T}^{(2)}$ the Fibonacci tree with root label 2. The trees are called Fibonacci trees for the following reason. Let $\#_n(\mathbb{T}^{(j)})$ denote the number of vertices in the n-th generation of the tree $\mathbb{T}^{(j)}$, the root being the first generation. Moreover, let f_n denote the n-th Fibonacci number starting with $f_1 = 1,\ f_2 = 1$. Then, one has $\#_n(\mathbb{T}^{(1)}) = f_n$ and $\#_n(\mathbb{T}^{(2)}) = f_{n+1}$.



Figure 1. Fibonacci tree $\mathbb{T}^{(1)}$ associated to substitution matrix $\left(\begin{smallmatrix} 0 & 1 \\ 1 & 1 \end{smallmatrix}\right)$. Vertices of label 1 are filled circles and vertices of label 2 are non-filled circles.

2010 *Mathematics Subject Classification.* Primary 82B44, Secondary 47B80, 60H25.

Key words and phrases. random Schrodinger operators, Anderson model, Fibonacci tree, extended states, absolutely continuous spectrum, localization.

For convenience, instead of considering these two trees, we will work with the forest $\mathbb{T} = \mathbb{T}^{(1)} \cup \mathbb{T}^{(2)}$ where the union has to be understood as disjoint union. By $d(x,y)$ we denote the graph distance of $x, y \in \mathbb{T}$, where $d(x,y) = \infty$ if x and y are in the different connected components.

A Fibonacci tree-strip is the cross product of $\mathbb{T}^{(1)}$ or $\mathbb{T}^{(2)}$ with a finite set $\mathcal{I} = \{1, \ldots, m\}$. On $\ell^2(\mathbb{T} \times \mathcal{I}) \cong \ell^2(\mathbb{T}, \mathbb{C}^m) = \{u : \mathbb{T} \to \mathbb{C}^m, \sum_{x \in \mathbb{T}} \|u(x)\|^2 < \infty\}$ which is also canonically equivalent to $\ell^2(\mathbb{T}^{(1)}, \mathbb{C}^m) \oplus \ell^2(\mathbb{T}^{(2)}, \mathbb{C}^m)$ and $\ell^2(\mathbb{T}) \otimes \mathbb{C}^m$, we define the random operators

$$(H_\lambda u)(x) = \left(\sum_{y : d(x,y) = 1} u(y) \right) + Au(x) + \lambda V(x) u(x) \,. \tag{1.1}$$

$A \in \mathrm{Sym}(m)$ represents the 'free vertical operator' and the matrices $V(x) \in \mathrm{Sym}(m)$ for $x \in \mathbb{T}$ are independent identically distributed random variables, distributed according to some probability measure ν on $\mathrm{Sym}(m)$ and scaled by the coupling constant λ. Here, $\mathrm{Sym}(m)$ denotes the set of real symmetric $m \times m$ matrices. These operators might be either thought of to model one particle on the product $\mathbb{T} \times \mathcal{I}$ or to model one particle on the Fibonacci tree with internal degrees of freedom and random hopping between these internal degrees, described by A and $V(x)$. Clearly, $H_\lambda = H_\lambda^{(1)} \oplus H_\lambda^{(2)}$, where $H_\lambda^{(q)}$ is the restriction of H_λ to $\ell^2(\mathbb{T}^{(q)}, \mathbb{C}^m)$ and can be seen as random Schrödinger operator on the tree strip $\mathbb{T}^{(q)} \times \mathcal{I}$.

If $\mathcal{I} = \mathbb{G}$ is a finite graph, then $\mathbb{T} \times \mathbb{G}$ can be interpreted as the product graph where $(x,k), (y,j) \in \mathbb{T} \times \mathbb{G}$ are connected by an edge, if either $x = y$ and $k, j \in \mathbb{G}$ are connected by an edge, or $k = j$ and $x, y \in \mathbb{T}$ are connected by an edge. If A is chosen to be the adjacency operator on $\mathbb{G}$, then H_0 is the adjacency operator on this product graph and H_λ is the Anderson model on this product graph.

For the Anderson model on $\mathbb{Z}^d$ or $\mathbb{R}^d$ in any dimension d, Anderson localization is proved at spectral edges and for high disorder [FS, FMSS, DLS, SW, CKM, DK, Kl2, AM, Aiz, Wa, Klo]. It is also known to hold for one dimensional [GMP, KuS, CKM] and quasi-one dimensional models like strips [Lac, KLS] and finite dimensional trees [Breu], except if built in symmetries prevent localization (e.g. [SS]).

In dimensions $d \geq 3$, it is expected to obtain some absolutely continuous (a.c.) spectrum for low disorder whereas for $d = 2$ one expects localization. These conjectures remain open problems. The existence of a.c. spectrum has only been proved for the Anderson model on trees and other tree-like graphs of infinite dimension with expoentially growing boundary. [Kl3, Kl4, Kl6, ASW, FHS, FHS2, Hal, AW, KLW, KLW2, FHH, KS, Sad]. This work adds another example to this list. It appears that the hyperbolic nature of such graphs leads to conservation of a.c. spectrum and ballistic dynamical behavior [Kl5, KS2, AW2] and these results should hold for much more general hyperbolic graphs. Therefore, it may be worth it to further geralize the results and identify the technical problems occuring in this process.

In my previous work [Sad] I considered random Schrödinger operators on certain tree-strips of finite cone type. If a tree has an assigned root 0 then the n-th generation is the set of vertices x with graph distance $d(x,0) = n$. The cone of descendants of x is then defined as the set of vertices y, where the shortest path to the root goes through x, i.e. the set of y such that $d(0,y) = d(0,x) + d(x,y)$.

The phrase 'finite cone type' refers to the fact, that for these trees, there are only finitely many different (isomorphy classes of) cones of descendants. This is also true for the Fibonacci trees. Cutting the connection to the root at a vertex of label p, $p = 1, 2$, leaves a cone of descendants isomorphic to $\mathbb{T}^{(p)}$, so there are only two different types of cones of descendants. However, the Fibonacci tree was not covered in [Sad]. One of the assumptions needed in [Sad] was that every vertex has at least 2 children which plays a significant role at various places. This means the trees could not have any line segment, that is a vertex not being the root which has only two neighbors. The Fibonacci tree is the simplest not-one dimensional tree of finite cone type which does not satisfy this assumption.

The main argument in [Sad] is adapted from [Kl3, KS] and uses a fixed point equation and the Implicit Function Theorem performed in some Banach spaces that are associated to supersymmetric functions. As we will see, for the Fibonacci tree this technique also works, but there are quite a few technical subtleties that are pointed out in this work. The Implicit Function Theorem is applied in a slightly different Banach space ($\mathcal{H} \times \mathcal{H}_\infty$ instead of $\mathcal{H}_\infty^2$). For the analysis of the Frechet derivative, compactness of a certain operator is needed which demands some additional work in this case (cf. Lemma 5.1). For the induction argument in Proposition 6.3 to obtain pure a.c. spectrum this Banach space is not good enough. One somehow needs to get back to the Banach space as used in [Sad] which is done through the argument in Proposition 6.1, especially (6.3) and (6.4) (cf. Remark 6.2). This relies on the identities mentioned in Appendix B and one needs some stronger assumption on the distribution of the matrix valued potential $V(x)$, namely it has to be compactly supported. For the Fibonacci tree one can explicitly calculate the spectrum for the adjacency operator (cf. Proposition 1.1), therefore the main Theorem here is less technical. Moreover, in [Sad, Theorem 1.2] some set of energies had to be excluded to get the almost sure a.c. spectrum. This set was given by a very technical condition which was shown to remove a nowhere dense set in a certain case. In Lemma 5.2 we show that this condition is never satisfied in the Fibonacci tree-strip case, hence we do not have to remove certain energies. The argument is based on the identity in Appendix A and works generally for trees of finite cone-type associated to a 2×2 substitution matrix S with negative determinant (cf. Remark 1.3).

Considering (1.1), let us remark that there is an orthogonal matrix $O \in \mathrm{O}(2)$ such that $O^\top A O$ is diagonal. Then $(\mathbf{1} \otimes O)$ is unitary and one obtains the equivalent family of operators

$$[(\mathbf{1} \otimes O)^* H_\lambda (\mathbf{1} \otimes O)]\, u(x) = \left(\sum_{y: d(x,y)=1} u(y) \right) + O^\top A O u(x) + \lambda O^\top V(x) O u(x) \;.$$

Hence, without loss of generality, we can assume that A is a diagonal matrix and we will do so in the proofs. In particular, the non-random operator H_0 is unitarily equivalent to a direct sum of shifted adjacency operators on $\mathbb{T}$, $H_0 = \boldsymbol{\Delta} \otimes \mathbf{1} + \mathbf{1} \otimes A \cong \bigoplus_{j=1}^m \Delta + a_j$ on $\ell^2(\mathbb{T}) \otimes \mathbb{C}^m$ where the a_j are the eigenvalues of A and Δ describes the adjacency operator on $\ell^2(\mathbb{T})$ given by

$$(\Delta v)(x) = \sum_{y: d(x,y)=1} v(y) \;, \qquad v \in \ell^2(\mathbb{T}) \;. \tag{1.2}$$

Our interest lies in the spectral type of H_λ which is determined by the matrix-valued spectral measures at the vertices of the forest $\mathbb{T}$. For $x \in \mathbb{T}$, $j \in \mathcal{I} = \{1,\ldots,m\}$ let $|x,j\rangle$ denote the element in $\ell^2(\mathbb{T},\mathbb{C}^m)$ satisfying $|x,j\rangle(y) = \delta_{x,y} e_j$ where e_j is the j-th canonical basis vector in $\mathbb{C}^m$. Moreover, for an operator H we denote by $\langle x,j|H|y,k\rangle$ the scalar product between $|x,j\rangle$ and $H|y,k\rangle$ with the convention that the scalar product is linear in the second and anti-linear in the first component. Then, for $x \in \mathbb{T}$ we define the random, positive matrix valued measure μ_x on $\mathbb{R}$ by

$$\int f(E)\, d\mu_x = [\, \langle x,j|f(H_\lambda)|x,k\rangle \,]_{j,k\in\mathcal{I}} \tag{1.3}$$

for all compactly supported, continuous functions f on $\mathbb{R}$. The roots of $\mathbb{T}^{(q)} \subset \mathbb{T}$, $q = 1,2$, will have a special role, therefore let us denote them by $0_q \in \mathbb{T}^{(q)}$.

In Section 2 we will see that:

Proposition 1.1. *The spectrum of the adjacency operator* Δ *on* $\mathbb{T}$ *is purely absolutely continuous and given by*

$$I = \left[-\frac{3}{2}\sqrt{3}\,,\, \frac{3}{2}\sqrt{3} \right] . \tag{1.4}$$

Therefore, letting $a_{\min} = a_1 \leq a_2 \leq \ldots \leq a_m = a_{\max}$ be the eigenvalues of A, one obtains that the the essential spectrum of H_0 is given by the union of the bands

$$\bigcup_{j=1}^{m} (I + a_j) = \left[-\frac{3}{2}\sqrt{3} + a_{\min}\,,\, \frac{3}{2}\sqrt{3} + a_{\max} \right] . \tag{1.5}$$

However, for the perturbation theory to work, the considered energies have to be in the intersection of these bands, and we need to exclude the centers of these bands. Therefore, let us define

$$I_A = \bigcap_{j=1}^{m} ((\mathring{I} + a_j) \setminus \{a_j\}) \;=\; \left(-\frac{3}{2}\sqrt{3} + a_{\max}\,,\, \frac{3}{2}\sqrt{3} + a_{\min} \right) \;\setminus \{a_1, a_2, \ldots, a_m\} . \tag{1.6}$$

Assumptions. *The following assumptions turn out to be crucial for the results.*

(V) *The distribution of* $V(x)$ *is compactly supported. As a consequence, all mixed finite moments of the random entries of the random matrix* $V(x)$ *exist and all partial derivatives of the Fourier transform*

$$h(M) := \mathbb{E}\left(e^{-i\mathrm{Tr}(MV(x))} \right) , \quad M \in \mathrm{Sym}(m) \tag{1.7}$$

exist to any order and are bounded.

(A) *Letting* $a_{\min} = a_1 \leq a_2 \leq \ldots \leq a_m = a_{\max}$ *be the eigenvalues of* A, *we assume*

$$a_{\max} - a_{\min} < 3\sqrt{3} , \tag{1.8}$$

This is equivalent to the set I_A *not being empty.*

Theorem 1.2. *Let the assumptions* (A) *and* (V) *be satisfied. There is an open neighborhood* U *of* $\{0\} \times I_A$ *in* $\mathbb{R}^2$ *such that the following holds:*

(i) *The spectrum of* H_λ *is almost surely purely absolutely continuous in* $U_\lambda = \{E : (\lambda, E) \in U\}$.

(ii) *For every $x \in \mathbb{T} \setminus \{0_1\}$, i.e. every vertex in $\mathbb{T}$ except the root of $\mathbb{T}^{(1)}$, the density of the absolutely continuous average spectral measure $\mathbb{E}(\mu_x)$ in U_λ depends continuously on $(\lambda, E) \in U$.*

(iii) *The density of $\mathbb{E}(\mu_{0_2})$ and $\mathbb{E}(\mu_{1_1})$ are positive definite in U_λ, where 1_1 is the child and only neighbor of the root $0_1 \in \mathbb{T}^{(1)}$ This implies that there is spectrum in U_λ with positive probability.*

Remark 1.3.

(i) *The set I_A here corresponds to $I_{A,S}$ in* [Sad]*. One of the difference to the previous work is that one needed to remove some energies and consider a very technical defined smaller set $\hat{I}_{A,S}$. This set occurred because the main argument is based on the Implicit Function Theorem and one needs a certain Frechet derivative to be invertible. Here, this Frechet derivative will always be invertible for all $E \in I_A$ by Lemma 5.2 which relies on the identity given in Proposition 2.1 This identity holds more general for the Green's functions on trees of finite cone type as shown in Appendix A. Using this identity and the same line of argument as in Lemma 5.2, one can actually show that for any 2×2 matrix S with negative determinant, the sets $I_{A,S}$ and $\hat{I}_{A,S}$ as defined in* [Sad] *coincide.*

(ii) *The root 0_1 of $\mathbb{T}^{(1)}$ is special as it is the only vertex of $\mathbb{T}$ which has only one neighbor. This is also the main reason that it has to be excluded in part (ii) of Theorem 1.2.*

The important objects we work with are the matrix Green's functions given by

$$G_\lambda^{[x]}(z) := \left[\, \langle x, j | (H - z)^{-1} | x, k \rangle \,\right]_{j,k \in \mathcal{I}} \in \mathbb{C}^{m \times m} \tag{1.9}$$

for $\mathrm{Im}(z) > 0$. The most important ingredient to obtain Theorem 1.2 is the following.

Theorem 1.4. *Under assumptions* (V) *and* (A) *there exists an open neighborhood U of $\{0\} \times I_A$ in $\mathbb{R}^2$ such that for all vertices $x \in \mathbb{T} \setminus \{0_1\}$ (i.e. all vertices except for the root of $\mathbb{T}^{(1)}$) the functions*

$$(\lambda, E, \eta) \mapsto \mathbb{E}\left(G_\lambda^{[x]}(E + i\eta) \right) ,$$
$$(\lambda, E, \eta) \mapsto \mathbb{E}\left(\left| G_\lambda^{[x]}(E + i\eta) \right|^2 \right) ,$$

defined for $\eta > 0$, have continuous extensions to $U \times [0, \infty)$.

Let us show now that Theorem 1.4 implies Theorem 1.2.

Proof of Theorem 1.2. Part (ii) follows immediately as $\mathbb{E}(G_\lambda^{[x]}(z))$ is the Stieltjes transform of $\mathbb{E}(\mu_x)$. For any compact interval $[a, b] \subset U_\lambda$ one has by Fatou's lemma

$$\mathbb{E}\Big(\liminf_{\mathrm{Im}(z) \downarrow 0} \int_a^b \mathrm{Tr}\big(\big| G_\lambda^{[x]}(z) \big|^2 \big)\, dE \Big) \le \liminf_{\mathrm{Im}(z) \downarrow 0} \int_a^b \mathbb{E}\Big(\mathrm{Tr}\big(\big| G_\lambda^{[x]}(z) \big|^2 \big) \Big)\, dE \, < \, \infty$$

Thus,

$$\liminf_{\mathrm{Im}(z) \downarrow 0} \int_a^b \mathrm{Tr}\left(\left| G_\lambda^{[x]}(z) \right|^2 \right) dE \, < \, \infty \quad \text{with probability one.} \tag{1.10}$$

As $G^{[x]}_\lambda(z)$ is the Stieltjes transform of μ_x this implies that, almost surely, μ_x is absolutely continuous with respect to the Lebesgue measure in $(a,b)\subset U_\lambda$ and the density is a matrix valued L^2 function for $x\in\mathbb{T}$, $x\neq 0_1$ (cf. [Kl6, Theorem 4.1] and [Kel, Theorem 2.6]). As there are countably many vertices $x\in\mathbb{T}$ and the open set U_λ can be obtained as countable union of intervals (a,b) satisfying $[a,b]\subset U_\lambda$, we obtain that with probability one, all measures μ_x ($x\neq 0_1$) are absolutely continuous in U_λ. As the vectors $\{|x,k\rangle\,:\,x\neq 0_1\}$ and $\{H_\lambda|1_1,k\rangle\}$ together span $\ell^2(\mathbb{T}\times\mathcal{I})$ this implies part (i). Part (iii) follows from the continuity in Theorem 1.4 around $\lambda=0$ for a possibly smaller neighborhood $U\supset\{0\}\times I_A$. $\square$

The paper is structured as follows. In Section 2 the spectrum of the adjacency operator on $\mathbb{T}$ and some crucial identities are derived. Then, in Section 3, we introduce the important Banach spaces that were also used in the previous works. In Section 4 we derive the fixed point equations and investigate the Frechet derivative in Section 5. Finally, in Section 6 we conclude to obtain Theorem 1.4. Appendices A and B state some general identities that are used along the way.

2. The unperturbed Green's functions

Similarly to above, we let $|x\rangle$ denote the function $|x\rangle(y)=\delta_{x,y}$ on $\ell^2(\mathbb{T})$ and for $\mathrm{Im}(z)>0$ we define the Green's functions

$$\Gamma^{(q)}_z := \langle 0_q|(\Delta - z)^{-1}|0_q\rangle \tag{2.1}$$

and the spectral measures ν_q at the roots 0_q of $\mathbb{T}^{(q)}$ for $q=1,2$ by

$$\int f(x)\,d\nu_q(x) \;=\; \langle 0_q|f(\Delta)|0_q\rangle\;. \tag{2.2}$$

Furthermore, we define

$$\Gamma^{(q)}_E := \lim_{\eta\downarrow 0}\Gamma^{(q)}_{E+i\eta} \tag{2.3}$$

for real energies E, if this limit exists as a limit in $\mathbb{C}$ (if the limit approaches infinity it is considered as non-existent). Furthermore, let us define the set $\hat{I}\subset\sigma(\Delta)$ by

$$\hat{I} := \left\{E\in\mathbb{R}\,:\,\Gamma^{(q)}_E=\lim_{\eta\downarrow 0}\Gamma^{(q)}_{E+i\eta}\ \text{ exists in } \mathbb{C} \text{ and }\ \mathrm{Im}(\Gamma^{(q)}_E)>0\ \text{ for } q=1,2\right\}\;. \tag{2.4}$$

Note that cutting the connections to the root and removing the root in the tree $\mathbb{T}^{(1)}$ we obtain $\mathbb{T}^{(2)}$ and doing the same in $\mathbb{T}^{(2)}$ we obtain two trees, one of which is $\mathbb{T}^{(1)}$ and the other $\mathbb{T}^{(2)}$. Therefore, the standard recursion for the Green's functions that can be obtained from the resolvent identity (cf. [FHS, KLW, Kel]) is given by

$$\Gamma^{(1)}_z = \frac{-1}{z+\Gamma^{(2)}_z}\;,\qquad \Gamma^{(2)}_z = \frac{-1}{z+\Gamma^{(1)}_z+\Gamma^{(2)}_z}\;. \tag{2.5}$$

From the first equation we see that in order for E to be in I, it is enough that the limit of $\lim_{\eta\downarrow 0}\Gamma^{(2)}_{E+i\eta}$ exists and has positive imaginary part.

By an hyperbolic contraction argument as in [FHS, KLW, Kel] one obtains that for $\mathrm{Im}(z)>0$, these equations together with the restriction $\mathrm{Im}(\Gamma^{(q)}_z)>0$, $q=1,2$; determine the Green's functions uniquely. By continuity, for $E=z\in\hat{I}$ these equations are also fulfilled and $\hat{I}$ is given by the set of real energies E, where the system

$$x = \frac{-1}{E+y}\;, \qquad y = \frac{-1}{E+x+y} \tag{2.6}$$

has a solution (x, y) with $\operatorname{Im}(x) > 0$ and $\operatorname{Im}(y) > 0$. From the first equation it is clear that $\operatorname{Im}(x) > 0$ if $\operatorname{Im}(y) > 0$.

After some algebra one obtains that $\hat{I}$ is therefore given by the set of energies $E \in \mathbb{R}$ where one obtains a solution y with positive imaginary part of

$$y^3 + 2Ey^2 + E^2 y + E = 0\;. \tag{2.7}$$

The discriminant of this polynomial in y is given by

$$D = 36E^4 - 32E^4 + 4E^6 - 4E^6 - 27E^2 = 4E^4 - 27E^2 = E^2(4E^2 - 27)\;. \tag{2.8}$$

The equation (2.7) has three real roots if $D \geq 0$ and one real and two complex-conjugate solutions if $D < 0$. Therefore,

$$\hat{I} = \left(-\frac{3\sqrt{3}}{2}\;;\; \frac{3\sqrt{3}}{2} \right) \setminus \{0\} \tag{2.9}$$

Moreover, for $E \in \hat{I}$, $\Gamma_E^{(2)}$ is given by the solution of (2.7) with positive imaginary part, which is

$$\begin{aligned} \Gamma_E^{(2)} = &-\frac{2E}{3} + \frac{1+i\sqrt{3}}{6}\sqrt[3]{\frac{1}{2}\left[27E - 2E^3 + \sqrt{27E^2(27-4E^2)}\right]} \\ &+ \frac{1-i\sqrt{3}}{6}\sqrt[3]{\frac{1}{2}\left[27E - 2E^3 - \sqrt{27E^2(27-4E^2)}\right]} \end{aligned} \tag{2.10}$$

For any z with $\operatorname{Im} z > 0$ we find from (2.5)

$$\operatorname{Im}\Gamma_z^{(2)} = \frac{\operatorname{Im}(z + \Gamma_z^{(1)} + \Gamma_z^{(2)})}{\left|z + \Gamma_z^{(1)} + \Gamma_z^{(2)}\right|^2} \leq \frac{1}{\operatorname{Im}(z + \Gamma_z^{(1)} + \Gamma_z^{(2)})} \leq \frac{1}{\operatorname{Im}\Gamma_z^{(2)}}$$

which implies the uniform bound $\operatorname{Im}\Gamma_z^{(2)} \leq 1$. Therefore, the spectral measure ν_2 is purely absolutely continuous and supported on $I = [-\frac{3}{2}\sqrt{3}, \frac{3}{2}\sqrt{3}]$, the closure of $\hat{I}$. The only case where $\operatorname{Im}\Gamma_{E+i\eta}^{(1)}$ can approach infinity is if $\Gamma_{E+i\eta}^{(2)}$ approaches $-E$. Plugging $y = -E$ into (2.7) gives $E = 0$. And indeed, at this particular energy, $\Gamma_{0+i\eta}^{(2)}$ approaches 0 and $\Gamma_{0+i\eta}^{(1)}$ approaches ∞ (for $\eta \downarrow 0$). Hence, for $E \notin I$ we have $E \neq 0$ and $\lim_{\eta\downarrow}\operatorname{Im}(\Gamma_{E+i\eta}^{(1)}) = 0$ by formula (2.5). Therefore, the support of ν_1 is also given by I.

An induction argument (induction over distance to root) as done in [Kel, Prop. 2.9] then shows, that for all other points $x \in \mathbb{T}$ the spectral measure at x is supported in I and hence I is equal to the spectrum of the adjacency operator Δ on $\mathbb{T}$. Moreover, this argument gives boundedness of the corresponding Green's functions for energies away from $E = 0$. So the only possible singular spectrum would be an eigenvalue with energy $E = 0$.

Assume such an eigenvector u exists. It satisfies $\sum_{y:d(x,y)=1} u(x) = 0$. For some point y of label 2, consider the child x of label 1 which has just one child y' of label 2. Then, $u(y') = -u(y)$. Iterating this argument, we see that u can not be in $\ell^2(\mathbb{T})$ unless $u(y) = 0$ for all y of label 2. Now, for a point x of label 1 consider the child y of label 2 and its children x' of label 1 and y' of label 2. Then

$u(x') = -u(x) - u(y') = -u(x)$. The same argument now shows $u(x) = 0$ for all x of label 1, if u is in $\ell^2(\mathbb{T})$. Hence, there is no eigenvector and 0 is not an eigenvalue. Therefore, the spectrum $\sigma(\Delta)$ is purely absolutely continuous and given by I and we just proved Proposition 1.1.

For our further analysis we summarize the important properties of $\Gamma^{(1)}_E$ and $\Gamma^{(2)}_E$ in the following proposition.

Proposition 2.1. *For* $E \in \hat{I} = (-\frac{3}{2}\sqrt{3}, \frac{3}{2}\sqrt{3}) \setminus \{0\}$ *the limits* $\Gamma^{(1)}_E = \lim_{\eta \to 0} \Gamma^{(1)}_{E+i\eta}$ *and* $\Gamma^{(2)}_E = \lim_{\eta\to 0} \Gamma^{(2)}_{E+i\eta}$ *exist and have a positive imaginary part. Moreover, for* $E \in \hat{I}$ *one has*

$$\left|\Gamma^{(2)}_E\right|^2 + \left|\Gamma^{(1)}_E \Gamma^{(2)}_E\right|^2 = 1 \,. \tag{2.11}$$

Proof. The first part follows from the formulas (2.5) and (2.10). Equation (2.11) can be rewritten as

$$\det\left[\begin{pmatrix} \left|\Gamma^{(1)}_E\right|^2 & 0 \\ 0 & \left|\Gamma^{(2)}_E\right|^2 \end{pmatrix}\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} - \mathbf{1}\right] = 0 \tag{2.12}$$

which is an identity shown for a more general case of trees of finite cone type in Proposition A.1 in Appendix A.

□

Remark 2.2. *The importance of the defined set* I_A *now comes from the following fact.*

$$I_A = \bigcap_{j=1}^{m} (\hat{I} + a_j) \tag{2.13}$$

Hence, if $E \in I_A$ *then* $E - a_j \in \hat{I}$ *for all* a_j *and we can use the proposition above for the shifted Green's functions that will appear in several formulas.*

3. Some Banach spaces

Here we briefly introduce the important Banach spaces and operators as in [KS, KS2, Sad]. All proofs and arguments are omitted as [Sad, Section 3] and uses the exact same notations and has the statements in more detail. For a supersymmetric background of these definitions see [Sad, Appendix B] or [KS].

Let $\mathcal{I} = \{1, \ldots, m\}$ and let $\mathfrak{P}(\mathcal{I}) = \{a : a \subset \mathcal{I}\}$ denote the set of all subsets of $\mathcal{I}$. Furthermore, let $\mathrm{Sym}^+(m)$ denote the set of real, symmetric $m \times m$ matrices M satisfying $M \geq 0$ in matrix sense, i.e. for all $v \in \mathbb{R}^m$ one has $v^\top M v \geq 0$. We define $\mathcal{P}$ to be the set of pairs $(\bar{a}, a)$ of subsets of $\mathcal{I}$ with the same cardinality,

$$\mathcal{P} := \{(\bar{a}, a) \,:\, \bar{a}, a \subset \mathcal{I},\, |\bar{a}| = |a|\} \,. \tag{3.1}$$

Moreover, let $n \in \mathbb{Z}$, $n \geq \frac{m}{2}$, and let $\bar{\mathbf{a}} = (\bar{a}_1, \ldots, \bar{a}_1)$, $\mathbf{a} = (a_1, \ldots, a_n) \in (\mathfrak{P}(\mathcal{I}))^n$ and define

$$\mathcal{P}^n := \{(\bar{\mathbf{a}}, \mathbf{a}) \in (\mathfrak{P}(\mathcal{I}))^n \times (\mathfrak{P}(\mathcal{I}))^n \,:\, (\bar{a}_l, a_l) \in \mathcal{P}\} \,. \tag{3.2}$$

For functions on $(\mathrm{Sym}^+(m))$ let $\partial_{j,k}$ denote the derivative with respect to the j,k- entry of M, $\partial_{j,k} f(M) = \frac{\partial}{\partial M_{j,k}} f(M)$ (by symmetry, $\partial_{j,k} = \partial_{k,j}$) and let $\tilde{\partial}_{j,k} = \frac{1}{2}(1 + \delta_{j,k})\partial_{j,k}$, where $\delta_{j,k}$ denotes the Kronecker delta symbol, $\delta_{j,k} = 0$ for $j \neq k$, and $\delta_{j,j} = 1$. For $(\bar{a}, a) \in \mathcal{P}$ with $a = \{k_1, \ldots, k_c\}$, $k_1 < k_2 \ldots < k_c$; $\bar{a} = \{\bar{k}_1, \ldots, \bar{k}_c\}$, $\bar{k}_1 < \bar{k}_2 < \ldots \bar{k}_c$, we define as in [KS, KS2]

$$\boldsymbol{\partial}_{\bar{a},a} := \begin{pmatrix} \tilde{\partial}_{\bar{k}_1,k_1} & \cdots & \tilde{\partial}_{\bar{k}_1,k_c} \\ \vdots & \ddots & \vdots \\ \tilde{\partial}_{\bar{k}_c,k_1} & \cdots & \tilde{\partial}_{\bar{k}_c,k_c} \end{pmatrix} . \tag{3.3}$$

Now set $D_{\emptyset,\emptyset}$ to be the identity operator and define

$$D_{\bar{a},a} := \det(\boldsymbol{\partial}_{\bar{a},a}) \quad \text{and} \quad D_{\bar{\mathbf{a}},\mathbf{a}} := \prod_{\ell=1}^{n} D_{\bar{a}_\ell,a_\ell} . \tag{3.4}$$

for $(\bar{a}, a) \in \mathcal{P}$ and $(\bar{\mathbf{a}}, \mathbf{a}) \in \mathcal{P}^n$. Something quite important is the following Leibniz-type rule. There is a function $(\bar{\mathbf{a}}, \mathbf{a}, \bar{\mathbf{b}}, \mathbf{b}, \bar{\mathbf{b}}', \mathbf{b}') \mapsto \operatorname{sgn}(\bar{\mathbf{a}}, \mathbf{a}, \bar{\mathbf{b}}, \mathbf{b}, \bar{\mathbf{b}}', \mathbf{b}') \in \{-1, 0, 1\}$ such that

$$D_{\bar{\mathbf{a}},\mathbf{a}}\,(fg) = \sum_{(\bar{\mathbf{b}},\mathbf{b}),(\bar{\mathbf{b}}',\mathbf{b}') \in \mathcal{P}^n} \operatorname{sgn}(\bar{\mathbf{a}}, \mathbf{a}, \bar{\mathbf{b}}, \mathbf{b}, \bar{\mathbf{b}}', \mathbf{b}') (D_{\bar{\mathbf{b}},\mathbf{b}}\, g)\, (D_{\bar{\mathbf{b}}',\mathbf{b}'}\, f) . \tag{3.5}$$

The exact expression of the function sgn above is explained in more detail in [Sad, Appendix B], in many cases one actually has $\operatorname{sgn}(\bar{\mathbf{a}}, \mathbf{a}, \bar{\mathbf{b}}, \mathbf{b}, \bar{\mathbf{b}}', \mathbf{b}') = 0$.

For $\boldsymbol{\varphi}, \boldsymbol{\varphi}' \in \mathbb{R}^{m \times 2n}$, $B \in \mathbb{C}^{m \times m}$ we define

$$\boldsymbol{\varphi} \cdot B\boldsymbol{\varphi}' := \operatorname{Tr}(\boldsymbol{\varphi}^\top B \boldsymbol{\varphi}') , \quad \boldsymbol{\varphi}^{\odot 2} := \boldsymbol{\varphi}\boldsymbol{\varphi}^\top \in \operatorname{Sym}^+(m) . \tag{3.6}$$

Let $C_n^\infty(\operatorname{Sym}^+(m))$ denote the set of smooth functions f on the interior of $\operatorname{Sym}^+(m)$ such that the functions $\boldsymbol{\varphi} \mapsto D_{\bar{\mathbf{a}},\mathbf{a}} f(\boldsymbol{\varphi}^{\odot 2})$ extend to C^∞ functions on $\mathbb{R}^{m \times 2n}$. (Since $2n \geq m$, $\boldsymbol{\varphi}^{\odot 2}$ is in the interior of $\operatorname{Sym}^+(m)$ if $\boldsymbol{\varphi}$ has full rank, hence $\boldsymbol{\varphi} \mapsto f(\boldsymbol{\varphi}^{\odot 2})$ is well defined for a dense open set in $\mathbb{R}^{m \times 2n}$. Moreover, let $\mathcal{S}_n(\operatorname{Sym}^+(m))$ denote the set of functions where $\boldsymbol{\varphi} \mapsto D_{\bar{\mathbf{a}},\mathbf{a}} f(\boldsymbol{\varphi}^{\odot 2})$ extends to a Schwartz function. A smooth function $\boldsymbol{\varphi} \mapsto g(\boldsymbol{\varphi})$ is a Schwartz function if for any polynomial function $p(\boldsymbol{\varphi})$ of the entries of $\boldsymbol{\varphi}$ and any combination of derivatives $D^\alpha = \prod_{j,k} \left(\frac{\partial}{\partial \boldsymbol{\varphi}_{j,k}} \right)^{\alpha_{j,k}}$ for a multi-index α, one has $\sup_{\boldsymbol{\varphi}} |p(\boldsymbol{\varphi}) D^\alpha g(\boldsymbol{\varphi})| < \infty$.

For $f \in \mathcal{S}_n(\operatorname{Sym}^+(m))$ we introduce the norms as in [Kl1, KSp, KS, KS2, Sad]

$$|||f|||_p^2 := \sum_{(\bar{\mathbf{a}},\mathbf{a}) \in \mathcal{P}^n} \left\| 2^{|\mathbf{a}|} D_{\bar{\mathbf{a}},\mathbf{a}}\, f\,(\boldsymbol{\varphi}^{\odot 2}) \right\|^2_{L^p(\mathbb{R}^{m \times 2n}, d^{2mn}\boldsymbol{\varphi})} . \tag{3.7}$$

where $|\mathbf{a}| = |a_1| + |a_2| + \ldots + |a_n|$. Now let $\widehat{\mathcal{H}}$ be the completion of $\mathcal{S}_n(\operatorname{Sym}^+(m))$ with respect to the norm $|||\cdot|||_2$, $\widehat{\mathcal{H}}$ is a Hilbert space. The Banach spaces $\widehat{\mathcal{H}}_p$, $p \in [1, \infty]$, are defined by

$$\widehat{\mathcal{H}}_p := \{ f \in \widehat{\mathcal{H}} \,:\, \|f\|_{\widehat{\mathcal{H}}_p} := |||f|||_2 + |||f|||_p < \infty \} . \tag{3.8}$$

So $f \in \widehat{\mathcal{H}}_p$ basically means that for all $(\bar{\mathbf{a}}, \mathbf{a}) \in \mathcal{P}^n$ the function $\boldsymbol{\varphi} \mapsto D_{\bar{\mathbf{a}},\mathbf{a}} f(\boldsymbol{\varphi}^{\odot 2})$ is an L^2 and L^p function of $\boldsymbol{\varphi}$.

For the same technical reasons as in [KS, Sad] we have to work on some specific closed subspaces.

Definition 3.1.

(i) *Let* $\operatorname{Sym}_{\mathbb{C}}(m)$ *denote the complex symmetric,* $m \times m$ *matrices.*

(ii) *For* $B \in \mathrm{Sym}_{\mathbb{C}}(m)$ *with strictly positive real part (i.e.,* $\operatorname{Re} B > 0$*), let* $\mathcal{PE}(B)$ *denote the vector space spanned by functions* $f \in C_n^\infty(\mathrm{Sym}^+(m))$ *of the form* $f(M) = p(M)\exp(-\mathrm{Tr}(MB))$*, where* $p(M)$ *is a polynomial in the entries of* $M \in \mathrm{Sym}^+(m)$*. Clearly,* $\mathcal{PE}(B) \subset \mathcal{S}_n(\mathrm{Sym}^+(m))$*.*

(iii) *Let* $\mathcal{PE}^{(0)}(B) = \{f \in \mathcal{PE}(B) : f(0) = 0\}$*.*

(iv) *Define* $\mathcal{PE}(m) \subset \mathcal{S}_n(\mathrm{Sym}^+(m))$ *as the smallest vector spaces containing all vector spaces* $\mathcal{PE}(B)$ *for all* $B \in \mathrm{Sym}_{\mathbb{C}}(m)$ *with* $\mathrm{Re}(B) > 0$*.*

(v) *For* $p \in [1, \infty]$ *let* $\mathcal{H}$ *and* $\mathcal{H}_p$ *be the closures of* $\mathcal{PE}(m)$ *in* $\widehat{\mathcal{H}}$ *and* $\widehat{\mathcal{H}}_p$*, respectively.*

On $\mathcal{S}_n(\mathrm{Sym}^+(m)) \otimes \mathcal{S}_n(\mathrm{Sym}^+(m))$ let us introduce the product norms

$$|||g|||_p^2 := \sum_{(\bar{\mathbf{a}},\mathbf{a}),(\bar{\mathbf{b}},\mathbf{b}) \in \mathcal{P}^n} \left\| 2^{|\mathbf{a}|+|\mathbf{b}|} D^{(+)}_{\bar{\mathbf{a}},\mathbf{a}} D^{(-)}_{\bar{\mathbf{b}},\mathbf{b}} g(\boldsymbol{\varphi}_+^{\odot 2}, \boldsymbol{\varphi}_-^{\odot 2}) \right\|^2_{L^p(\mathbb{R}^{4mn}, d^{2mn}\boldsymbol{\varphi}_+ d^{2mn}\boldsymbol{\varphi}_-)} , \tag{3.9}$$

where $D^{(\pm)}_{\bar{\mathbf{a}},\mathbf{a}}$ denotes the operator $D_{\bar{\mathbf{a}},\mathbf{a}}$ with respect to the entry $\boldsymbol{\varphi}_\pm^{\odot 2}$. Using these norms we define the following spaces.

Definition 3.2.

(i) *Let* $\widehat{\mathcal{K}} = \hat{\mathcal{H}} \otimes \hat{\mathcal{H}}$ *which is the completion of* $\mathcal{S}_n(\mathrm{Sym}^+(m)) \otimes \mathcal{S}_n(\mathrm{Sym}^+(m))$ *with respect to the norm* $|||\cdot|||_2$*.*

(ii) *Let* $\widehat{\mathcal{K}}_p := \{g \in \hat{\mathcal{K}} : |||g|||_2 + |||g|||_p < \infty\}$ *.*

(iii) *Let* $\mathcal{K}$ *be the closure of* $\mathcal{PE}(m) \otimes \mathcal{PE}(m)$ *in* $\widehat{\mathcal{K}}$*.*

(iv) *For* $p \in [1, \infty]$ *let* $\mathcal{K}_p$ *be the closure of* $\mathcal{PE}(m) \otimes \mathcal{PE}(m)$ *in* $\widehat{\mathcal{K}}_p$*, respectively.*

Lemma 3.3 in [Sad] sates:

Lemma 3.3. *Given any complex symmetric* $m \times m$ *matrix* B *with* $\operatorname{Re} B > 0$ *and* $p \in [1, \infty)$*,* $\mathcal{H}$ *and* $\mathcal{H}_p$ *are the closures of* $\mathcal{PE}(B)$ *in* $\widehat{\mathcal{H}}$ *and* $\widehat{\mathcal{H}}_p$*, respectively. Similarly,* $\mathcal{PE}(B) \otimes \mathcal{PE}(C)$ *is dense in* $\mathcal{K}$ *and* $\mathcal{K}_p$ *for any symmetric* $m \times m$ *matrices* B*,* C *with* $\mathrm{Re}(B) > 0$ *and* $\mathrm{Re}(C) > 0$*.*

As in [KS, KS2, Sad] let us introduce the supersymmetric Fourier transform T acting on $\mathcal{S}_n(\mathrm{Sym}^+(m))$ by

$$Tf(\boldsymbol{\varphi}'^{\odot 2}) = \frac{(-1)^{mn}}{\pi^{mn}} \int e^{\pm i \boldsymbol{\varphi} \cdot \boldsymbol{\varphi}'} D_{\boldsymbol{\mathcal{I}},\boldsymbol{\mathcal{I}}} f(\boldsymbol{\varphi}^{\odot 2}) \, d^{2mn}\boldsymbol{\varphi} \tag{3.10}$$

For this definition it is important that $2n \geq m$, because this insures that the map $\boldsymbol{\varphi} \mapsto \boldsymbol{\varphi}^{\odot 2}$ from $\mathbb{R}^{m \times 2n}$ to $\mathrm{Sym}^+(m)$ is surjective and hence $Tf \in \mathcal{S}_n(\mathrm{Sym}^+(m))$ is well defined. Note, that the right hand side does not depend on the sign of $\pm\boldsymbol{\varphi} \cdot \boldsymbol{\varphi}'$ in the exponent. One finds (cf. [Sad, eq. (B.22)], [KS, eq. (2.37)])

$$D_{\bar{\mathbf{a}},\mathbf{a}}(Tf) = \tfrac{2^{mn}}{4^{|\mathbf{a}|}} \operatorname{sgn}(\mathbf{a}, \bar{\mathbf{a}}) \, \mathcal{F}(D_{\mathbf{a}^c, \bar{\mathbf{a}}^c} f) \quad \text{for all} \quad (\bar{\mathbf{a}}, \mathbf{a}) \in \mathcal{P}^n , \tag{3.11}$$

where $\mathcal{F}f(\boldsymbol{\varphi}^{\odot 2})$ denotes the Fourier transform w.r.t. to $\boldsymbol{\varphi}$ in $\mathbb{R}^{2mn}$ and $\operatorname{sgn}(\mathbf{a}, \bar{\mathbf{a}}) \in \{-1, 1\}$ is some sign.

Another key identity is the following equation which is derived in [Sad, eq. (B.26)]. Let B be a symmetric matrix and $\mathrm{Im}(B) > 0$. Then,

$$T(e^{i \mathrm{Tr}(B \boldsymbol{\varphi}^{\odot 2})}) = e^{-\frac{i}{4} \mathrm{Tr}(B^{-1} \boldsymbol{\varphi}^{\odot 2})} \tag{3.12}$$

Lemma 2.6 in [KS] states:

Lemma 3.4. (i) *T is unitary on* $\widehat{\mathcal{H}}$ *and* $\mathcal{H}$*.*

(ii) *T is a bounded operator from $\widehat{\mathcal{H}}_1$ to $\widehat{\mathcal{H}}_\infty$, as well as from $\mathcal{H}_1$ to $\mathcal{H}_\infty$.*

The operator $\mathcal{T} := T \otimes T$ is given by

$$\begin{aligned} &\mathcal{T} g((\boldsymbol{\varphi}'^{\odot 2}_{+}, \boldsymbol{\varphi}'^{\odot 2}_{-}) \qquad (3.13) \\ &\quad = \frac{1}{\pi^{2mn}} \int e^{\pm i(\boldsymbol{\varphi}'_+ \cdot \boldsymbol{\varphi}_+ \pm \boldsymbol{\varphi}'_- \cdot \boldsymbol{\varphi}_-)} D^{+}_{\boldsymbol{\mathcal{I}},\boldsymbol{\mathcal{I}}} D^{-}_{\boldsymbol{\mathcal{I}},\boldsymbol{\mathcal{I}}}\, g(\boldsymbol{\varphi}_+^{\odot 2}, \boldsymbol{\varphi}_-^{\odot 2})\, d^{2mn}\boldsymbol{\varphi}_+\, d^{2mn}\boldsymbol{\varphi}_- \end{aligned}$$

where $D^{\pm}_{\boldsymbol{\mathcal{I}},\boldsymbol{\mathcal{I}}}$ denotes the operator $D_{\boldsymbol{\mathcal{I}},\boldsymbol{\mathcal{I}}}$ with respect to the entry $\boldsymbol{\varphi}_{\pm}^{\odot 2}$. $\mathcal{T}$ is unitary on $\widehat{\mathcal{K}}$, $\mathcal{K}$ and $\mathcal{K}^{(0)}$ and it defines a bounded linear map from $\widehat{\mathcal{K}}_1$ to $\widehat{\mathcal{K}}_\infty$, from $\mathcal{K}_1$ to $\mathcal{K}_\infty$ and from $\mathcal{K}_1^{(0)}$ to $\mathcal{K}_\infty^{(0)}$.

4. FIXED POINT EQUATIONS

In this and the following sections let the assumption (V) hold. For two neighboring sites x, $y \in \mathbb{T}^{(q)}$ let $\mathbb{T}^{(x|y)}$ denote the rooted tree with root x obtained by removing the branch from x going through y in $\mathbb{T}^{(q)}$. Furthermore, let $H_\lambda^{(x|y)}$ denote the operator H_λ restricted to $\mathbb{T}^{(x|y)}$ with Dirichlet boundary conditions and similar to (1.9) let

$$G_\lambda^{(x|y)}(z) := \left[\langle x, j | (H_\lambda^{(x|y)} - z)^{-1} | x, k\rangle\right]_{j,k} \in \mathbb{C}^{m\times m} \,. \qquad (4.1)$$

For simplicity, we will denote the Green's functions at the roots by

$$G_\lambda^{(q)}(z) := G_\lambda^{[0_q]}(z) = \left[\langle 0_q, j | (H_\lambda - z)^{-1} | 0_q, k\rangle\right]_{j,k} \in \mathbb{C}^{m\times m} \,. \qquad (4.2)$$

As in [Sad] define for any $z = E + i\eta$ with $\eta > 0$, and label $q = 1, 2$, $\zeta_{\lambda,z}^{(q)}$, $\zeta_{\lambda,z}^{(x|y)}$ on $\mathrm{Sym}^+(m)$ and $\xi_{\lambda,z}^{(q)}$, $\xi_{\lambda,z}^{(x|y)}$ on $\mathrm{Sym}^+(m) \times \mathrm{Sym}^+(m)$ by

$$\zeta_{\lambda,z}^{(q)}(\boldsymbol{\varphi}^{\odot 2}) := \mathbb{E}\left(\mathrm{e}^{\frac{i}{2}\mathrm{Tr}(G_\lambda^{(q)}(z)\,\boldsymbol{\varphi}^{\odot 2})}\right) \,, \qquad (4.3)$$

$$\zeta_{\lambda,z}^{(x|y)}(\boldsymbol{\varphi}^{\odot 2}) := \mathbb{E}\left(\mathrm{e}^{\frac{i}{2}\mathrm{Tr}(G_\lambda^{(x|y)}(z)\,\boldsymbol{\varphi}^{\odot 2})}\right) \,, \qquad (4.4)$$

$$\xi_{\lambda,z}^{(q)}(\boldsymbol{\varphi}_+^{\odot 2}\,,\, \boldsymbol{\varphi}_-^{\odot 2}) := \mathbb{E}\left(e^{\frac{i}{2}\mathrm{Tr}\left(G_\lambda^{(q)}(z)\,\boldsymbol{\varphi}_+^{\odot 2} - \overline{G_\lambda^{(q)}(z)}\,\boldsymbol{\varphi}_-^{\odot 2}\right)}\right) \,, \qquad (4.5)$$

$$\xi_{\lambda,z}^{(x|y)}(\boldsymbol{\varphi}_+^{\odot 2}\,,\, \boldsymbol{\varphi}_-^{\odot 2}) := \mathbb{E}\left(e^{\frac{i}{2}\mathrm{Tr}\left(G_\lambda^{(x|y)}(z)\,\boldsymbol{\varphi}_+^{\odot 2} - \overline{G_\lambda^{(x|y)}(z)}\,\boldsymbol{\varphi}_-^{\odot 2}\right)}\right) \,. \qquad (4.6)$$

As in the previous works, let us introduce the operators

$$B_{\lambda,z} = \mathcal{M}\left(e^{i\mathrm{Tr}(\frac{1}{2}(z-A)\boldsymbol{\varphi}^{\odot 2})}\ h(\tfrac{1}{2}\lambda\boldsymbol{\varphi}^{\odot 2})\right) \,, \qquad (4.7)$$

$$\mathcal{B}_{\lambda,z} = \mathcal{M}\left(e^{i\mathrm{Tr}(\frac{1}{2}(z-A)\boldsymbol{\varphi}_+^{\odot 2} - (\frac{1}{2}(\bar{z}-A)\boldsymbol{\varphi}_-^{\odot 2}))}\ h(\tfrac{1}{2}\lambda(\boldsymbol{\varphi}_+^{\odot 2} - \boldsymbol{\varphi}_-^{\odot 2}))\ \right) \qquad (4.8)$$

where $\mathcal{M}(g)$ denotes the multiplication operator which acts on function by multiplication with g. Moreover, h denotes the Fourier transform of the distribution of $V(x)$ as given in (1.7).

Let N_p denote the set of neighbors of the root 0_p, i.e. $N_p = \{x : d(x, 0_p) = 1\}$. The well-known recursion relation for the Green's function on trees is given by

$$G_\lambda^{(p)}(z) = -\left(z - \lambda V(0_p) - A + \sum_{x\in N_p} G_\lambda^{(x|0_p)}(z)\right)^{-1} \qquad (4.9)$$

which by (3.12) is equivalent to

$$e^{\frac{i}{2}\,\mathrm{Tr}\left(G_\lambda^{(p)}(z)\boldsymbol{\varphi}^{\odot 2}\right)} = T\mathcal{M}(e^{i\mathrm{Tr}((z-\lambda\frac{1}{2}V(x)-\frac{1}{2}A)\boldsymbol{\varphi}^{\odot 2})})\left(\prod_{x\in N_p} e^{\frac{i}{2}\mathrm{Tr}(G_\lambda^{(x|0_p)}(z)\boldsymbol{\varphi}^{\odot 2})}\right)\,. \tag{4.10}$$

Using the independence of the matrix potentials, this immediately implies

$$\zeta^{(1)}_{\lambda,z} = TB_{\lambda,z}\zeta^{(2)}_{\lambda,z}\,,\quad \zeta^{(2)}_{\lambda,z} = TB_{\lambda,z}\left(\zeta^{(1)}_{\lambda,z}\zeta^{(2)}_{\lambda,z}\right)\,. \tag{4.11}$$

Analogously, one also obtains

$$\xi^{(1)}_{\lambda,z} = \mathcal{TB}_{\lambda,z}\xi^{(2)}_{\lambda,z}\,,\quad \xi^{(2)}_{\lambda,z} = \mathcal{TB}_{\lambda,z}\left(\xi^{(1)}_{\lambda,z}\xi^{(2)}_{\lambda,z}\right)\,. \tag{4.12}$$

Recall that we assume without loss of generality that $A = \mathrm{diag}(a_1,\ldots,a_m)$ is diagonal. Therefore, in the free case, $\lambda = 0$, one obtains

$$\zeta^{(q)}_{0,z}(\boldsymbol{\varphi}^{\odot 2}) = e^{\frac{i}{2}\mathrm{Tr}(A^{(q)}_z\boldsymbol{\varphi}^{\odot 2})}\ ,\ \text{ where }\ A^{(q)}_z := \mathrm{diag}(\Gamma^{(q)}_{z-a_1},\Gamma^{(q)}_{z-a_2},\ldots,\Gamma^{(q)}_{z-a_m})\,. \tag{4.13}$$

By Proposition 2.1, for $E\in I_A$ the point-wise limits

$$\zeta^{(q)}_{0,E}(\boldsymbol{\varphi}^{\odot 2}) := \lim_{\eta\downarrow 0}\zeta^{(q)}_{0,E+i\eta}(\boldsymbol{\varphi}^{\odot 2}) = e^{\frac{i}{2}\mathrm{Tr}(A^{(q)}_E\,\boldsymbol{\varphi}^{\odot 2})}, \tag{4.14}$$

$$\xi^{(q)}_{0,E}(\boldsymbol{\varphi}^{\odot 2}_+,\boldsymbol{\varphi}^{\odot 2}_-) := \lim_{\eta\downarrow 0}\xi^{(q)}_{0,E+i\eta}(\boldsymbol{\varphi}^{\odot 2}_+,\boldsymbol{\varphi}^{\odot 2}_-) = e^{\frac{i}{2}\mathrm{Tr}(A^{(q)}_E\,\boldsymbol{\varphi}^{\odot 2}_+ - \bar{A}^{(q)}_E\,\boldsymbol{\varphi}^{\odot 2}_-)} \tag{4.15}$$

exist for $q = 1,2$, where

$$A^{(q)}_E = \lim_{\eta\downarrow 0} A^{(q)}_{E+i\eta}\,,\qquad [A^{(q)}_E]_{jk} = \delta_{jk}\,\Gamma^{(q)}_{E-a_j} \tag{4.16}$$

are diagonal $m\times m$ matrices with strictly positive imaginary part. For simplified notations, let

$$\vec{\zeta}_{\lambda,z} = \begin{pmatrix}\zeta^{(1)}_{\lambda,z}\\ \zeta^{(2)}_{\lambda,z}\end{pmatrix}\quad\text{and}\quad \vec{\xi}_{\lambda,z} = \begin{pmatrix}\xi^{(1)}_{\lambda,z}\\ \xi^{(2)}_{\lambda,z}\end{pmatrix}\,. \tag{4.17}$$

Proposition 4.1. *We have:*

(i) *For $\eta = \mathrm{Im}\, z \geq 0$ the operator $B_{\lambda,z}$ is a bounded operator on on $\mathcal{H}$ and $\mathcal{H}_1$. The map*

$$F\,:\,(\lambda,E,\eta,f_1,f_2)\mapsto TB_{\lambda,E+i\eta}\begin{pmatrix}f_2\\ f_1f_2\end{pmatrix} \tag{4.18}$$

is a continuous map from $\mathbb{R}\times\mathbb{R}\times[0,\infty)\times\mathcal{H}\times\mathcal{H}_\infty$ to $\mathcal{H}\times\mathcal{H}_\infty$.

(ii) *$\vec{\zeta}_{\lambda,z}\in\mathcal{H}^2_\infty\subset\mathcal{H}\times\mathcal{H}_\infty$ for all $\lambda\in\mathbb{R}$ and $z = E+i\eta$ with $\eta>0$. The map $(\lambda,E,\eta)\to\vec{\zeta}_{\lambda,E+i\eta}$ is continuous from $\mathbb{R}\times\mathbb{R}\times(0,\infty)$ to $\mathcal{H}\times\mathcal{H}_\infty$.*

(iii) *If $E\in I_A$, then $\vec{\zeta}_{0,E}\in(\mathcal{PE}(m))^2\subset\mathcal{H}\times\mathcal{H}_\infty$ and*

$$\lim_{\eta\downarrow 0}\vec{\zeta}_{0,E+i\eta}\ =\ \vec{\zeta}_{0,E}\quad\text{in }\mathcal{H}\times\mathcal{H}_\infty\,. \tag{4.19}$$

(iv) *The equality (4.11) can be rewritten as a fixed point equation in $\mathcal{H}\times\mathcal{H}_\infty$:*

$$\vec{\zeta}_{\lambda,z}\ =\ F(\lambda,E,\eta,\vec{\zeta}_{\lambda,z}) \tag{4.20}$$

valid for all $\lambda\in\mathbb{R}$ and $z = E+i\eta$ with $\eta>0$, and also valid for $\lambda = 0$ and $z = E$ with $E\in I_A$.

Proof. (i) As mentioned in [Sad, Proposition 4.1] and proved in [KS, Proposition 3.2], $B_{\lambda,z}$ is a bounded operator on $\mathcal{H}_1$ and $(\lambda, E, \eta, f) \mapsto B_{\lambda,z} f$ is a continuous map from $\mathbb{R} \times \mathbb{R} \times [0, \infty) \times \mathcal{H}_1$ to $\mathcal{H}_1$. The main ingredient is Dominated Convergence. Completely analogue one can prove the same statement replacing $\mathcal{H}_1$ by $\mathcal{H}$. Assumption (V) is important for this observation, as $B_{\lambda,z}$ includes the Fourier transform of the distribution of $V(x)$ (cf. Definitions (1.7) and (4.7)).

Using Hölder's inequality we find that $(f_1, f_2) \mapsto (f_2, f_1 f_2)$ is a continuous map from $\mathcal{H} \times \mathcal{H}_\infty$ to $\mathcal{H} \times \mathcal{H}_1$. These facts together with Lemma 3.4 (ii) immediately imply (i).

Parts (ii) - (iv) are proved exactly the same way as [Sad, Proposition 4.1,(ii)-(iv)]. and one could replace $\mathcal{H} \times \mathcal{H}_\infty$ by $\mathcal{H}_\infty^2$. □

One also obtains the analogue results for the function $\vec{\xi}_{\lambda,z}$.

Proposition 4.2.

(i) *For* $\eta = \operatorname{Im} z \geq 0$ *the operator* $\mathcal{B}_{\lambda,z}$ *is a bounded operator on* $\mathcal{K}$ *and* $\mathcal{K}_1$. *Furthermore, the map*

$$Q \,:\, (\lambda, E, \eta, g_1, g_2) \mapsto \mathcal{T}\mathcal{B}_{\lambda,E+i\eta} \begin{pmatrix} g_2 \\ g_1 g_2 \end{pmatrix} \tag{4.21}$$

is a continuous map from $\mathbb{R} \times \mathbb{R} \times [0, \infty) \times \mathcal{K} \times \mathcal{K}_\infty$ *to* $\mathcal{K} \times \mathcal{K}_\infty$.

(ii) $\vec{\xi}_{\lambda,z} \in \mathcal{K}_\infty^2 \subset \mathcal{K} \times \mathcal{K}_\infty$ *for all* $\lambda \in \mathbb{R}$ *and* $z = E + i\eta$ *with* $\eta > 0$. *The map* $(\lambda, E, \eta) \to \vec{\xi}_{\lambda,E+i\eta}$ *is continuous from* $\mathbb{R} \times \mathbb{R} \times (0, \infty)$ *to* $\mathcal{K} \times \mathcal{K}_\infty$.

(iii) *If* $E \in I_A$, *then* $\vec{\xi}_{0,E} \in \mathcal{K} \times \mathcal{K}_\infty$ *and*

$$\lim_{\eta \downarrow 0} \vec{\xi}_{0,E+i\eta} \;=\; \vec{\xi}_{0,E} \quad \text{in } \mathcal{K} \times \mathcal{K}_\infty \;. \tag{4.22}$$

(iv) *The equality* (4.12) *can be rewritten as a fixed point equation in* $\mathcal{K} \times \mathcal{K}_\infty$*:*

$$\vec{\xi}_{\lambda,z} \;=\; Q(\lambda, E, \eta, \vec{\xi}_{\lambda,z}) \;, \tag{4.23}$$

valid for all $\lambda \in \mathbb{R}$ *and* $z = E + i\eta$ *with* $\eta > 0$, *and also valid for* $\lambda = 0$ *and* $z = E$ *with* $E \in I_A$.

Remark 4.3. *One of the differences between this work and* [Sad] *is that for fixed* (λ, E, η) *the maps* F *and* Q *are operators on* $\mathcal{H} \times \mathcal{H}_\infty$ *and* $\mathcal{K} \times \mathcal{K}_\infty$, *respectively, but not on* $\mathcal{H}_\infty^2$ *or* $\mathcal{K}_\infty^2$. *We can not use the space* $\mathcal{H}_\infty^2$ *as for* $\eta = 0$ *and* $f_2 \in \mathcal{H}_\infty$ *we can not show that* $B_{\lambda,E} f_2$ *is in* $\mathcal{H}_1$ *so that* T *would map it back to* $\mathcal{H}_\infty$. *We can only say that one lands in* $\mathcal{H}$ *after applying* T. *This in turn comes from the fact that vertices of label* 1 *do have only one child in the Fibonacci trees and therefore we do not have a product of more than one function after* $B_{\lambda,z}$ *in the first entry. Recall that in* [Sad] *every vertex had to have at least two children. This would cause some slight problem for Proposition 6.3, however it can be resolved through the statements* (6.3) *and* (6.4) *in Proposition 6.1 (cf. Remark 6.2).*

The other difference is, that we do not need the smaller spaces $\mathcal{H}_\infty^{(0)}$, $\mathcal{K}_\infty^{(0)}$ *introduced in* [Sad] *to avoid some further assumption (but one could work in the spaces* $\mathcal{H}^{(0)} \times \mathcal{H}_\infty^{(0)}$, $\mathcal{K}^{(0)} \times \mathcal{K}_\infty^{(0)}$ *if one wanted to).*

5. Spectrum of Frechet derivatives

At some point when calculating the spectrum of the Frechet derivative, it will be important that a certain power is a compact operator. As we need a more general case here compared to previous work, we give the compactness argument in detail in the following lemma.

Lemma 5.1.

(i) *Let h_1, h_2 be exponentially decaying, continuous functions on $\mathbb{R}^k$, let $\mathcal{F}$ denote the Fourier transform on $\mathbb{R}^k$ and let $B \in \mathrm{Sym}(k) \cap \mathrm{GL}(k,\mathbb{R})$. Then,*

$$\mathcal{M}(h_1)\mathcal{F}\mathcal{M}(h_2) \quad \textit{and} \quad \mathcal{M}(h_1)\mathcal{F}^*\mathcal{M}(e^{\frac{1}{2}ix\cdot Bx})\mathcal{F}\mathcal{M}(h_2) \tag{5.1}$$

are compact operators from $L^2(\mathbb{R}^k)$ to $L^p(\mathbb{R}^k)$ for any $p \in [1,\infty]$.

(ii) *For $f_1, f_2 \in \mathcal{PE}(m)$ and $A \in \mathrm{Sym}(m) \cap \mathrm{GL}(m)$ the operators*

$$\mathcal{M}(f_1)T\mathcal{M}(f_2)\,, \quad \textit{and} \quad \mathcal{M}(f_1)T\mathcal{M}(e^{i\mathrm{Tr}(A\boldsymbol{\varphi}^{\odot 2})})T\mathcal{M}(f_2) \tag{5.2}$$

are compact operators from $\mathcal{H}_p$ to $\mathcal{H}_q$ for any $p, q \in [1,\infty]$. (Note that the case where one Banach space is $\mathcal{H}$ is included as $\mathcal{H} = \mathcal{H}_2$ as a set, only the norm differs technically by a factor 2).

(iii) *For $g_1, g_2 \in \mathcal{PE}(m) \otimes \mathcal{PE}(m)$, $A_+, A_- \in \mathrm{Sym}(m) \cap \mathrm{GL}(m)$, the operators*

$$\mathcal{M}(g_1)\mathcal{T}\mathcal{M}(g_2)\,, \quad \textit{and} \quad \mathcal{M}(g_1)\mathcal{T}\mathcal{M}(e^{i\mathrm{Tr}(A_+\boldsymbol{\varphi}_+^{\odot 2} + A_-\boldsymbol{\varphi}_-^{\odot 2})})\mathcal{T}\mathcal{M}(g_2) \tag{5.3}$$

are compact operators from any $\mathcal{K}_p$ to any $\mathcal{K}_q$ for any $p, q \in [1,\infty]$.

Proof. For (i) let us first assume that h_1 and h_2 are compactly supported and consider $\mathcal{M}(h_1)\mathcal{F}\mathcal{M}(h_2)$. Let K be the compact support of h_2. There exists a constant C_K such that for all $x \in K$ and all $y \in \mathbb{R}^k$ we have $|e^{ix\cdot y} - e^{ix\cdot y'}| \leq C_K|y - y'|$. Therefore

$$\begin{aligned} |(\mathcal{F}\mathcal{M}(h_1)f)(y) - (\mathcal{F}\mathcal{M}(h_1)f)(y')| &\leq (2\pi)^{-k/2} C_K |y - y'|\, \|h_2 f\|_1 \\ &\leq (2\pi)^{-k/2} C_K \|h_2\|_2\, \|f\|_2\, |y - y'| \end{aligned}$$

and hence $\mathcal{M}(h_1)\mathcal{F}\mathcal{M}(h_2)$ maps a L^2 bounded sequence of functions into a sequence of equi-continuous functions, supported on the compact support of h_1. By the theorem of Arzela Ascoli we obtain a convergent subsequence in L^∞ and hence in any L^p norm.

If h_1 and h_2 are continuous and exponentially decaying, then we can approach them in $\|\cdot\|_\infty$ norm by compactly supported continuous functions $h_{1,n}$, $h_{2,n}$. Then $\mathcal{M}(h_{1,n})\mathcal{F}\mathcal{M}(h_{2,n})$ approaches $\mathcal{M}(h_1)\mathcal{F}\mathcal{M}(h_2)$ in $L^2 \to L^p$ operator norm for any $p \in [1,\infty]$ (here, consider $\mathcal{M}(h_1)$ as map from L^2 to L^1, $\mathcal{F}$ as map from L^1 to L^∞ and $\mathcal{M}(h_1)$ as map from L^∞ to L^p).

For the second operator in (5.1) note that by Lemma B.2 in Appendix B one finds $\mathcal{M}(h_1)\mathcal{F}^*\mathcal{M}(e^{ix\cdot Bx})\mathcal{F}\mathcal{M}(h_2) = \mathcal{S}_B\mathcal{M}(\hat{h}_1)\mathcal{F}\mathcal{M}(\hat{h}_2)$ where $\hat{h}_1$ and $\hat{h}_2$ are exponentially decaying functions and $\mathcal{S}_B f(x) = f(B^{-1}x)$. Therefore, by the previous statement, this defines a compact operator from L^2 to any L^p, $p \in [1,\infty]$.

Using (3.11) and the Leibniz-rule (3.5) the statements (ii) and (iii) immediately follow from (i). Starting with an $\mathcal{H}$ bounded (respectively, $\mathcal{K}$-bounded) sequence one can subsequently construct a subsequence converging in all norms involved in the definition of $\mathcal{H}_p$, $\mathcal{K}_p$ respectively. $\square$

It is now time to introduce some more notations to properly describe the spectrum of the Frechet derivatives. These notations were also used in [Sad]. By $\Delta(m, \mathbb{Z}_+)$ we denote the set of upper triangular matrices with non-negative integer entries. For $J \in \Delta(m, \mathbb{Z}_+)$ and $E \in I_A$ we define

$$\theta^{(q)}_{J,E} := \prod_{\substack{j,k \in \{1,\dots,m\} \\ j \le k}} \left[\Gamma^{(q)}_{E-a_j} \Gamma^{(q)}_{E-a_k} \right]^{J_{jk}} \in \mathbb{C}\,,\ q = 1, 2\,, \quad \text{and} \tag{5.4}$$

$$\boldsymbol{\theta}_{J,E} := \operatorname{diag}(\theta^{(1)}_{J,E}, \theta^{(2)}_{J,E}) = \begin{pmatrix} \theta^{(1)}_{J,E} & 0 \\ 0 & \theta^{(2)}_{J,E} \end{pmatrix} \tag{5.5}$$

With the help of these matrices we will express the spectrum of the important Frechet derivatives. The following result will ensure that we can use the Implicit Function Theorem.

Lemma 5.2. *For $E \in I_A$ and any $J, J' \in \Delta(m, \mathbb{Z}_+)$ we find*

$$\det(\boldsymbol{\theta}_{J,E} \boldsymbol{\theta}^*_{J',E} S - \mathbf{1}) \neq 0. \tag{5.6}$$

This means, the matrices $\boldsymbol{\theta}_{J,E}\boldsymbol{\theta}^*_{J',E} S$ do not have an eigenvalue 1.

Proof. For $J, J' \in \Delta(m, \mathbb{Z}_+)$ and $E \in I_A$ define

$$f(J, J', E) = \det(\boldsymbol{\theta}_{J,E}\boldsymbol{\theta}^*_{J',E} S - \mathbf{1}) = 1 - \theta^{(2)}_{J,E} \left(\theta^{(2)}_{J',E}\right)^* - \theta^{(1)}_{J,E} \left(\theta^{(1)}_{J',E}\right)^* \theta^{(2)}_{J,E} \left(\theta^{(2)}_{J',E}\right)^* . \tag{5.7}$$

For $J = J' = 0$ we have $f(0, 0, E) = \det(S - \mathbf{1}) = -1 \neq 0$. Now let $\|J\|_1$ denote the norm given by the sum of the absolute values of all entries of J. Next, we consider the case $\|J\|_1 + \|J'\|_1 = 1$, i.e. one of these matrices is zero and the other has one entry. Both cases are completely analogous so let us just consider $J' = 0$, $\|J\|_1 = 1$. Then by (5.4) one has

$$\theta^{(q)}_{J,E} = \Gamma^{(q)}_{E-b_1} \Gamma^{(q)}_{E-b_2} \tag{5.8}$$

for some $b_1, b_2 \in \{a_1, \dots, a_m\}$. Using (2.11) in Proposition 2.1 and the Cauchy-Schwarz inequality we find

$$\begin{aligned} \left|\theta^{(2)}_{J,E}\right| + \left|\theta^{(1)}_{J,E}\theta^{(2)}_{J,E}\right| &= \left|\Gamma^{(2)}_{E-b_1}\right| \left|\Gamma^{(2)}_{E-b_2}\right| + \left|\Gamma^{(1)}_{E-b_1}\Gamma^{(2)}_{E-b_1}\right| \left|\Gamma^{(1)}_{E-b_2}\Gamma^{(2)}_{E-b_2}\right| \\ &\le \prod_{k=1}^{2} \sqrt{\left|\Gamma^{(2)}_{E-b_k}\right|^2 + \left|\Gamma^{(1)}_{E-b_k}\Gamma^{(2)}_{E-b_k}\right|^2} = 1 \end{aligned} \tag{5.9}$$

Since $\theta^{(2)}_{J,E}$ is the product of two factors with positive imaginary part, it can not be a positive real number, hence $|f(J,0,E)| > 1 - |\theta^{(2)}_J| - |\theta^{(1)}_{J,1}\theta^{(2)}_{J,E}| \ge 0$, so $f(J, 0, E)$ can not be zero in this case.

Finally, consider $\|J\|_1 + \|J'\|_1 \ge 2$. We may assume without loss of generality, that $J \neq 0$. Then

$$\theta^{(q)}_{J,E} \left(\theta^{(q)}_{J',E}\right)^* = \Gamma^{(q)}_{E-b_1} \Gamma^{(q)}_{E-b_2} \cdot X^{(q)} \tag{5.10}$$

where $X^{(q)}$ itself is a product of an even number of factors (at least 2) $\Gamma^{(q)}_{E-b}$ or complex conjugates. By (2.11) and the fact that for $E \in I_A$ and $b \in \{a_1, \dots, a_m\}$ none of the imaginary parts of $\Gamma^{(q)}_{E-b}$ can be zero, we find $|\Gamma^{(2)}_{E-b}| < 1$ and $|\Gamma^{(1)}_{E-b}\Gamma^{(2)}_{E-b}| < 1$

leading to $|X^{(2)}| < 1$ and $|X^{(1)}X^{(2)}| < 1$. Using this and Cauchy-Schwartz as in (5.8) we find

$$\left|\theta^{(2)}_{J,E}\left(\theta^{(2)}_{J',E}\right)^*\right| + \left|\theta^{(1)}_{J,E}\left(\theta^{(1)}_{J',E}\right)^*\theta^{(2)}_{J,E}\left(\theta^{(2)}_{J',E}\right)^*\right| < 1 \tag{5.11}$$

which immediately implies $|f(J,J',E)| > 0$. Hence, in any case, $f(J,J',E)$ will not be zero. □

The following lemma corresponds to [Sad, Lemma 5.1].

Lemma 5.3.

(i) *The map F as in (4.18) is continuous and Frechet-differentiable w.r.t. $\vec{f} = (f_1, f_2) \in \mathcal{H} \times \mathcal{H}_\infty$. For $\vec{f} \in \mathcal{H}^2_\infty$ the Frechet derivative $F_{\vec{f}}$ extends naturally to a bounded operator on $\mathcal{H}^2$ which we will also denote as $F_{\vec{f}}$*

(ii) *For $E \in I_A$ let $C_E = F_{\vec{f}}(0, E, 0, \vec{\zeta}_{0,E})$, then C_E^4 is a compact operator on $\mathcal{H} \times \mathcal{H}_\infty$ and $\mathcal{H}^2$.*

(iii) *The spectrum of C_E as an operator on the Hilbert space $\mathcal{H}^2$ is given by the eigenvalues of the matrices $\boldsymbol{\theta}_{J,E}S$ for $J \in \Delta(m, \mathbb{Z}_+)$ and the accumulation point 0. This means, denoting the spectrum of C_E on $\mathcal{H}^2$ by $\sigma_{\mathcal{H}^2}$ one obtains*

$$\sigma_{\mathcal{H}^2}(C_E) = \bigcup_{J\in\Delta(m,\mathbb{Z}_+)} \sigma(\boldsymbol{\theta}_{J,E}S) \,\cup\, \{0\} \tag{5.12}$$

where $\boldsymbol{\theta}_{J,E}$ are the matrices as defined in (5.5). In particular, by Lemma 5.2 one has for $E \in I_A$

$$1 \notin \sigma_{\mathcal{H}^2}(C_E) \tag{5.13}$$

(iv) *The spectrum of C_E as an operator on $\mathcal{H} \times \mathcal{H}_\infty$, denoted by $\sigma_{\mathcal{H}\times\mathcal{H}_\infty}(C_E)$, is the same as its spectrum as an operator on $\mathcal{H}^2$,*

$$\sigma_{\mathcal{H}\times\mathcal{H}_\infty}(C_E) = \sigma_{\mathcal{H}^2}(C_E). \tag{5.14}$$

Proof. (i) The derivative $F_{\vec{f}}$ can be written as a matrix of operators and we obtain formally

$$F_{\vec{f}} = \begin{pmatrix} 0 & TB_{\lambda,z} \\ TB_{\lambda,z}\mathcal{M}(f_2) & TB_{\lambda,z}\mathcal{M}(f_1) \end{pmatrix}. \tag{5.15}$$

For $f_1, g_1 \in \mathcal{H}$ and $f_2, g_2 \in \mathcal{H}_\infty$ one finds $B_{\lambda,z}f_1g_2$, $B_{\lambda,z}f_2g_1 \in \mathcal{H}_1$ and by Lemma 3.4 $F_{\vec{f}}$ defines a bounded linear operator on $\mathcal{H} \times \mathcal{H}_\infty$. Similarly, if $f_1, f_2 \in \mathcal{H}_\infty$, then it is also a bounded linear operator on $\mathcal{H}^2$.

To get (ii) note that $\vec{\zeta}_{0,E} \in (\mathcal{PE}(m))^2$, and $B_{0,E} = \mathcal{M}(e^{\frac{i}{2}\mathrm{Tr}((E-A)\boldsymbol{\varphi}^{\odot 2})})$. For simplicity, let us simply write B for $B_{0,E}$, M_1 for $\mathcal{M}(\zeta^{(1)}_{0,E})$ and M_2 for $\mathcal{M}(\zeta^{(2)}_{0,E})$, then

$$C_E^2 = \begin{pmatrix} TBTBM_2 & TBTBM_1 \\ TBM_1TBM_2 & TBM_2TB + TBM_1TBM_1 \end{pmatrix}. \tag{5.16}$$

Squaring this matrix we get that each summand in the entries of C_E^4 contains some piece of the form M_iTBTBM_j or M_iTBM_j which are compact operators by Lemma 5.1.

Part (iii) is the complete same calculation as in [Sad, Lemma 5.1 (iii)] giving the eigenvalues of C_E. At this point it is important that C_E has a compact power, as it assures that except for 0, the whole spectrum of C_E consists of eigenvalues. As a by product one also obtains that the eigenfunctions are in $\mathcal{H}^2_\infty$.

Part (iv) follows by the same argument as [Sad, Lemma 5.1 (iv)], $\sigma_{\mathcal{H}\times\mathcal{H}_\infty}(C_E) \subset \sigma_{\mathcal{H}^2}(C_E)$ by compactness of C_E^4 in $\mathcal{H} \times \mathcal{H}_\infty$. Equality follows as one finds eigenfunctions corresponding to the eigenvalues of $\boldsymbol{\theta}_{J,E} S$ in $\mathcal{H}_\infty^2$. ☐

Analogue results hold for the map Q and are proved the same way.

Lemma 5.4.

(i) *The map Q is Frechet-differentiable w.r.t. $\vec{g} \in \mathcal{K} \times \mathcal{K}_\infty$. The derivative $Q_{\vec{g}}$ is a bounded linear operator on $\mathcal{K} \times \mathcal{K}_\infty$ and for $\vec{g} \in \mathcal{K}_\infty^2$ it extends naturally to a bounded operator on $\mathcal{K}^2$.*

(ii) *For $E \in I_A$ let $\mathcal{C}_E = Q_{\vec{g}}(0, E, 0, \vec{\xi}_{0,E})$, then $\mathcal{C}_E^4$ is a compact operator on $\mathcal{K} \times \mathcal{K}_\infty$ and $\mathcal{K}^2$.*

(iii) *The spectrum of $\mathcal{C}_E$ as an operator on the Hilbert space $\mathcal{K}^2$ is given by the eigenvalues of the matrices $\boldsymbol{\theta}_{J,E}\boldsymbol{\theta}^*_{J',E}S$ and the accumulation point 0, i.e. denoting the spectrum of $\mathcal{C}_E$ on $\mathcal{K}^2$ by $\sigma_{\mathcal{K}^2}(\mathcal{C}_E)$ one finds*

$$\sigma_{\mathcal{K}^2}(\mathcal{C}_E) = \bigcup_{J,J' \in \Delta(m,\mathbb{Z}_+)} \sigma(\boldsymbol{\theta}_{J,E}\boldsymbol{\theta}^*_{J',E}S) \cup \{0\} \tag{5.17}$$

where $\boldsymbol{\theta}_{J,E}, \boldsymbol{\theta}_{J',E}$ are the matrices as defined in (5.5). By Lemma 5.2 we find

$$1 \notin \sigma_{\mathcal{K}^2}(\mathcal{C}_E)\ . \tag{5.18}$$

(iv) *The spectrum of $\mathcal{C}_E$ as an operator on $\mathcal{K} \times \mathcal{K}_\infty$ is the same as its spectrum as an operator on $\mathcal{K}^2$:*

$$\sigma_{\mathcal{K}\times\mathcal{K}_\infty}(\mathcal{C}_E) = \sigma_{\mathcal{K}^2}(\mathcal{C}_E). \tag{5.19}$$

Proof. The proof is completely analogous to the one for Lemma 5.3. For (i) and (ii) note that the Frechet derivative of Q is given by

$$Q_{\vec{g}} = \begin{pmatrix} 0 & \mathcal{T}\mathcal{B}_{\lambda,z} \\ \mathcal{T}\mathcal{B}_{\lambda,z}\mathcal{M}(g_2) & \mathcal{T}\mathcal{B}_{\lambda,z}\mathcal{M}(g_1) \end{pmatrix}\ . \tag{5.20}$$

Part (iii) is completely analogue to [Sad, Lemma 5.2 (iii)] and (iv) follows by the same arguments as in Lemma 5.3. ☐

6. Conclusions

The most important ingredient for the proof of Theorem 1.4 is the following.

Proposition 6.1. *There exists an open set U with $\{0\} \times I_A \subset U$, such that the maps*

$$(\lambda, E, \eta) \in U \times (0,\infty) \ \mapsto\ \vec{\zeta}_{\lambda,E+i\eta} = (\zeta^{(1)}_{\lambda,E+i\eta}, \zeta^{(2)}_{\lambda,E+i\eta}) \in \mathcal{H} \times \mathcal{H}_\infty \tag{6.1}$$

$$(\lambda, E, \eta) \in U \times (0,\infty) \ \mapsto\ \vec{\xi}_{\lambda,E+i\eta} = (\xi^{(1)}_{\lambda,E+i\eta}, \xi^{(2)}_{\lambda,E+i\eta}) \in \mathcal{K} \times \mathcal{K}_\infty \tag{6.2}$$

have continuous extensions to $U \times [0,\infty)$ satisfying (4.20) and (4.23). For these, extensions, $\zeta^{(1)}_{\lambda,E} \in \mathcal{H}_\infty$, $\xi^{(1)}_{\lambda,E} \in \mathcal{K}_\infty$. Moreover, for $1 \le p < \infty$ the maps

$$(\lambda, E, \eta, f) \in U \times (0,\infty) \times \mathcal{H}_p \ \mapsto\ \zeta^{(1)}_{\lambda,E+i\eta} f \ \in \mathcal{H}_p \tag{6.3}$$

$$(\lambda, E, \eta, g) \in U \times (0,\infty) \times \mathcal{K}_p \ \mapsto\ \xi^{(1)}_{\lambda,E+i\eta} g \ \in \mathcal{K}_p \tag{6.4}$$

also extend continuously to $U \times [0,\infty)$.

Proof. By Lemma 5.3 and Lemma 5.4 we can use the Implicit Function Theorem on Banach Spaces as stated in [Kl6, Appendix B] for the functions $\hat{F}(\lambda, E, \eta, \vec{f}) = F(\lambda, E, \eta, \vec{f}) - \vec{f}$ and $\hat{Q}(\lambda, E, \eta, \vec{g}) = Q(\lambda, E, \eta, \vec{g}) - \vec{g}$ at the points $(0, E, 0, \vec{\zeta}_{0,E})$ and $(0, E, 0, \vec{\xi}_{0,E})$ with $E \in I_A$. Uniqueness of the continuous implicit function and the continuity properties of $\vec{\zeta}_{\lambda,E+i\eta}$ and $\vec{\xi}_{\lambda,E+i\eta}$ as stated in Proposition 4.1 and Proposition 4.2 give the first two continuous extensions.

Note that by (4.20), one has

$$\zeta^{(1)}_{\lambda,z} = TB_{\lambda,z}TB_{\lambda,z}(\zeta^{(1)}_{\lambda,z}\zeta^{(2)}_{\lambda,z}) \tag{6.5}$$

where $z = E + i\eta$. Defining

$$\tilde{F}(\lambda, E, \eta, V) = T\mathcal{M}(e^{\frac{i}{2}\mathrm{Tr}((E-A-\lambda V)\mathbf{\Phi}^{\odot 2})})TB_{\lambda,E+i\eta}(\zeta^{(1)}_{\lambda,E+i\eta}\zeta^{(2)}_{\lambda,E+i\eta})\,.$$

this means $\zeta^{(1)}_{\lambda,z} = \mathbb{E}(\tilde{F}(\lambda, E, \eta, V(x))$ where $\mathbb{E}$ refers to the expectation over the distribution of the potential $V(x)$. Also note that by (3.5) and Hölder's inequality, the product $B_{\lambda,z}\zeta^{(1)}_{\lambda,z}\zeta^{(2)}_{\lambda,z}$ extends continuously as a map from $(\lambda, E, \eta) \in U \times [0, \infty)$ to $\mathcal{H}_1$. By assumption (V) and possibly shrinking the open set U, we can assure that for $(\lambda, E) \in U$ the inverses $(E + i\eta - A - \lambda V(x))^{-1}$ exist almost surely and are uniformly bounded for $V(x)$ and (λ, E) varying in a compact subset of U. Let $\lambda_n \to \lambda, E_n \to E, \eta_n \to \eta$ (i.e. $z_n = E_n + i\eta_n \to z = E + i\eta$) and $f_n \to f$ in $\mathcal{H}_p$. Using Corollary B.3 in Appendix B, the Leibniz rule (3.5) and (3.11) one obtains that for large n, $\zeta^{(1)}_{\lambda_n,z_n} = \mathbb{E}\tilde{F}(\lambda_n, E_n, \eta_n, V(x))$ is uniformly bounded in $\mathcal{H}_\infty$. Moreover, $D_{\bar{\mathbf{a}},\mathbf{a}}\zeta^{(1)}_{\lambda_n,z_n}$ converges point-wise to $D_{\bar{\mathbf{a}},\mathbf{a}}\zeta^{(1)}_{\lambda,z} = \mathbb{E}\tilde{F}(\lambda, E, \eta, V(x))$ (as a functions of $\boldsymbol{\varphi}^{\odot 2}$) for any $(\bar{\mathbf{a}}, \mathbf{a}) \in \mathcal{P}^n$.

By the uniform $\mathcal{H}_\infty$ bound and the Leibniz rule (3.5), $|\!|\!|\zeta^{(1)}_{\lambda_n,z_n} f_n - \zeta^{(1)}_{\lambda_n,z_n} f|\!|\!|_p$ goes to zero. Using (3.5) and Dominated Convergence, $|\!|\!|\zeta^{(1)}_{\lambda_n,z_n} f - \zeta^{(1)}_{\lambda,z} f|\!|\!|_p$ goes to zero as well. Hence, $\zeta^{(1)}_{\lambda_n,z_n} f_n \to \zeta^{(1)}_{\lambda,z} f$ in $\mathcal{H}_p$.

The proof for the map $(\lambda, E, \eta, g) \mapsto \xi^{(1)}_{\lambda,E+i\eta}g$ is completely analogue. □

Remark 6.2. *We can not show that the functions* $(\lambda, E, \eta) \mapsto \zeta^{(1)}_{\lambda,E+i\eta}$, *and* $\xi^{(1)}_{\lambda,E+i\eta}$ *extend continuously as maps from* $U \times (0, \infty)$ *to* $\mathcal{H}_\infty$ *or* $\mathcal{K}_\infty$, *respectively. Convergence in the norm* $|\!|\!| \cdot |\!|\!|_\infty$ *and* $|\!|\!|\!| . |\!|\!|\!|_\infty$ *is a problem if we let* $\eta \to 0$. *For this reason, we can not perform the Implicit Function Theorem using the space* $\mathcal{H}^2_\infty$ *or* $\mathcal{K}^2_\infty$ *as in* [Sad]. *But the continuous extensions for the maps* (6.3) *and* (6.4) *allow to complete the argument in the same way as in* [Sad].

Proposition 6.3. *There is an open set* U, $\{0\} \times I_A \subset U$ *such that one has the following:*

(i) *Let* 1_1 *be the only child and only neighbor of* 0_1, *the root of* $\mathbb{T}^{(1)}$, *then the map*

$$(\lambda, E, \eta, f) \in \mathbb{R} \times \mathbb{R} \times (0, \infty) \times \mathcal{H} \;\mapsto\; \zeta^{(0_1|1_1)}_{\lambda,z} f \in \mathcal{H} \tag{6.6}$$

extends continuously on $(\lambda, E, \eta, f) \in U \times [0, \infty) \times \mathcal{H}$.

(ii) *For all* $x, y \in \mathbb{T}$ *with* $d(x, y) = 1$ *and* $x \neq 0_1$ *the maps*

$$(\lambda, E, \eta) \in \mathbb{R} \times \mathbb{R} \times (0, \infty) \;\mapsto\; \zeta^{(x|y)}_{\lambda,E+i\eta} \in \mathcal{H}\,, \tag{6.7}$$

$$(\lambda, E, \eta) \in \mathbb{R} \times \mathbb{R} \times (0, \infty) \;\mapsto\; \xi^{(x|y)}_{\lambda,E+i\eta} \in \mathcal{K}\,, \tag{6.8}$$

have continuous extensions to $(\lambda, E, \eta) \in U \times [0, \infty)$.

Proof. We let $z = E + i\eta$. As the graph $\mathbb{T}^{(0_1|1_1)}$ contains only the point 0_1 one finds $\zeta_{\lambda,z}^{(0_1|1_1)}(\boldsymbol{\varphi}^{\odot 2}) = \mathbb{E}(e^{\frac{i}{2}\mathrm{Tr}((z-A-\lambda V(0_1))^{-1}\boldsymbol{\varphi}^{\odot 2})})$. As in the argument above, by assumption (V) the inverse in the exponential is locally uniformly bounded for small λ. Together with Dominated Convergence, this shows claim (i).

If $d(x,y) = 1$ and x is a child of y (i.e. $d(x,0_q) = 1 + d(y,0_q)$), then $\mathbb{T}^{(x|y)}$ is equivalent to $\mathbb{T}^{(p)}$ if the label of x is p, $l(x) = p$, and therefore $\zeta_{\lambda,z}^{(x|y)} = \zeta_{\lambda,z}^{(p)}$. Hence, by Proposition 6.1, $(\lambda, z) \mapsto \zeta_{\lambda,z}^{(x|y)}$ extends continuously as a function mapping to $\mathcal{H}$.

Now let x be the parent of y. We will prove the continuous extension by induction over $d(x,0_q)$ for $x \in \mathbb{T}^{(q)}$, where the induction starts with $x = 0_2$ on the tree $\mathbb{T}^{(2)}$ and $x = 1_1$, the only child of 0_1 on the tree $\mathbb{T}^{(1)}$. Note that the recursion relations (4.9) and (4.10) can be applied at any vertex in a tree. Hence, if $N(x)$ are the neighbors of x, $y \in N(x)$, then one obtains

$$\zeta_{\lambda,z}^{(x|y)} = TB_{\lambda,z} \prod_{y' \in N(x), y' \neq y} \zeta_{\lambda,z}^{(y'|x)}\,, \quad \xi_{\lambda,z}^{(x|y)} = \mathcal{TB}_{\lambda,z} \prod_{y' \in N(x), y' \neq y} \xi_{\lambda,z}^{(y'|x)}\,.$$

Let y, y' be the children of 0_2, then $(\lambda, z) \mapsto \zeta_{\lambda,z}^{(0_2|y)} = TB_{\lambda,z}\zeta_{\lambda,z}^{(y'|0_2)}$ extends continuously as a function mapping to $\mathcal{H}$ because $\zeta_{\lambda,z}^{(y'|0_2)}$ does. If y, y' are the children of 1_1, then by (6.6), $(\lambda, z) \mapsto \zeta_{\lambda,z}^{(1_1|y)} = TB_{\lambda,z}\zeta_{\lambda,z}^{(0_1|1_1)}\zeta_{\lambda,z}^{(y'|1_1)}$ extends continuously as a function mapping to $\mathcal{H}$ as well.

For the induction argument let y be a child of x and x be a child of $\tilde{x}$ and assume that $(\lambda, z) \mapsto \zeta_{\lambda,z}^{(\tilde{x}|x)} \in \mathcal{H}$ extends continuously. If x has label 1 then $\zeta_{\lambda,z}^{(x|y)} = TB_{\lambda,z}\zeta_{\lambda,z}^{(\tilde{x}|x)}$ and if x has label 2, then $\zeta_{\lambda,z}^{(x|y)} = TB_{\lambda,z}\zeta_{\lambda,z}^{(y'|x)}\zeta_{\lambda,z}^{(\tilde{x}|x)}$ where y' is the other child of x and $\zeta_{\lambda,z}^{(y'|x)}$ is equal to $\zeta_{\lambda,z}^{(1)}$ or $\zeta_{\lambda,z}^{(2)}$. In any of these cases, Proposition 6.1 gives that $(\lambda, z) \mapsto \zeta_{\lambda,z}^{(x|y)} \in \mathcal{H}$ extends continuously.

With similar arguments one obtains the statements for the functions $\xi_{\lambda,z}^{(x|y)}$. □

As shown in previous works [KS, Sad], the expected matrix valued Green's functions can be obtained by

$$\mathbb{E}\big(G_\lambda^{[x]}(z)\big) = -i \int \mathbf{D} B_{\lambda,z} \left(\prod_{y: d(x,y)=1} \zeta_{\lambda,z}^{(y|x)}(\boldsymbol{\varphi}^{\odot 2}) \right) d^{2mn}\boldsymbol{\varphi}\,, \tag{6.9}$$

$$\mathbb{E}\left(\left| G_\lambda^{[x]}(z) \right|^2 \right) = \int \mathbf{D}^{(-)}\mathbf{D}^{(+)} \mathcal{B}_{\lambda,z} \left(\prod_{y: d(x,y)=1} \xi_{\lambda,z}^{(y|x)}(\boldsymbol{\varphi}_+^{\odot 2}, \boldsymbol{\varphi}_-^{\odot 2}) \right) d^{2mn}\boldsymbol{\varphi}_+\, d^{2mn}\boldsymbol{\varphi}_-\,, \tag{6.10}$$

where $\mathbf{D}$ is an $m \times m$ matrix of differential operators with entries

$$\mathbf{D} := \left(\tfrac{(-1)^{mn+j+k}}{2\pi^{mn}}\, D_{\mathcal{I},\mathcal{I}}^{n-1}\, D_{\mathcal{I}\setminus\{k\},\mathcal{I}\setminus\{j\}} \right)_{j,k \in \mathcal{I}}\,. \tag{6.11}$$

with $D_{\mathcal{I},\mathcal{I}}$ and $D_{\mathcal{I}\setminus\{k\},\mathcal{I}\setminus\{j\}}$ as in (3.4). $\mathbf{D}^{(\pm)}$ represent the matrix-operator $\mathbf{D}$ acting with respect to $\boldsymbol{\varphi}_\pm^{\odot 2}$ and $\mathbf{D}^{(-)}\mathbf{D}^{(+)}$ has to be understood as a matrix product.

For $x = 0_2$ the product in (6.9) is $\zeta_{\lambda,z}^{(1)}\zeta_{\lambda,z}^{(2)}$, if x is not a root and has label 1, then the product is of the form $\zeta_{\lambda,z}^{(2)}\zeta_{\lambda,z}^{(y|x)}$ and if the label of x is 2, it is of the form

$\zeta^{(1)}_{\lambda,z}\zeta^{(2)}_{\lambda,z}\zeta^{(y|x)}_{\lambda,z}$. We have seen that $\zeta^{(2)}_{\lambda,z} \in \mathcal{H}_\infty$ and $\zeta^{(y|x)}_{\lambda,z} \in \mathcal{H}$ extend continuously as functions of $(\lambda, E, \eta) \in U \times [0,\infty)$ with $z = E + i\eta$. Using the Leibniz rule (3.5), Hölder's inequality and (6.3), $(\lambda, z) \mapsto \zeta^{(1)}_{\lambda,z}\zeta^{(2)}_{\lambda,z} \in \mathcal{H}_1$, $(\lambda, z) \mapsto \zeta^{(2)}_{\lambda,z}\zeta^{(y|x)}_{\lambda,z} \in \mathcal{H}_1$ and $(\lambda, z) \mapsto \zeta^{(1)}_{\lambda,z}\zeta^{(2)}_{\lambda,z}\zeta^{(y|x)}_{\lambda,z} \in \mathcal{H}_1$ extend continuously as functions mapping to $\mathcal{H}_1$. Similarly statements hold for ζ replaced by ξ and $\mathcal{H}, \mathcal{H}_p$ replaced by $\mathcal{K}$, $\mathcal{K}_p$. Thus, any of the functions integrated over in (6.9) or (6.10) are in $L^1(d^{2mn}\boldsymbol{\varphi})$ and $L^1(d^{2mn}\boldsymbol{\varphi}_+ d^{2mn}\boldsymbol{\varphi}_-)$, respectively, continuously depending on (λ, E, η) and Theorem 1.4 follows.

Appendix A. An identity for the unperturbed Green's functions on trees of finite cone type

Associated to an $s \times s$ substitution matrix $S \in \mathrm{Mat}(s, \mathbb{Z}_+)$ with non-negative integer entries are the following s rooted trees of finite cone type, denoted by $\mathbb{T}^{(r)}, r = 1, \ldots, s$. Each vertex has a label, the root of the tree $\mathbb{T}^{(r)}$ has label r. Any vertex of label p has $S_{p,q}$ children of label q. Denoting by Δ the adjacency operator on the forest $\bigcup_r \mathbb{T}^{(r)}$ and by $0_r \in \mathbb{T}^{(r)}$ the root of the tree $\mathbb{T}^{(r)}$, we define for $\mathrm{Im}(z) > 0$ the Green's functions

$$\Gamma^{(r)}_z = \langle 0_r \,|\, (\Delta - z)^{-1} \,|\, 0_r \rangle \,.$$

We define the set

$$\Sigma = \{E \in \mathbb{R} \,:\, \Gamma^{(r)}_E := \lim_{\eta \downarrow 0} \Gamma^{(r)}_{E+i\eta} \text{ exists for all } r \text{ and} \,\mathrm{Im}(\Gamma^{(q)}_E) > 0 \text{ for some } q\}$$

Proposition A.1. *Let* $\Gamma_E = \mathrm{diag}(\Gamma^{(1)}_E, \ldots, \Gamma^{(s)}_E)$ *denote the diagonal* $s \times s$ *matrix with the Green's functions along the diagonal. Then one has for* $E \in \Sigma$

$$\det(|\Gamma_E|^2 S - \mathbf{1}) \,=\, 0\,. \tag{A.1}$$

Proof. The recursion relation for the Green's functions is given by

$$\Gamma^{(p)}_E = -\left(E + \sum_{q=1}^{s} S_{p,q}\Gamma^{(q)}_E\right)^{-1} \,.$$

Multiplying by $(\Gamma^{(p)}_E)^*$ and some algebra leads to

$$\left|\Gamma^{(p)}_E\right|^2 \sum_{q=1}^{s} S_{p,q}\Gamma^{(q)}_E = -E\left|\Gamma^{(p)}_E\right|^2 - \left(\Gamma^{(p)}_E\right)^*$$

Taking imaginary parts gives

$$\left|\Gamma^{(p)}_E\right|^2 \sum_{q=1}^{s} S_{p,q}\,\mathrm{Im}(\Gamma^{(q)}_E) = \mathrm{Im}(\Gamma^{(p)}_E)$$

Defining the vector $\vec{\Gamma}_E = (\Gamma^{(1)}_E, \ldots, \Gamma^{(s)}_E)^\top$ this equation can be read as

$$|\Gamma|^2 S\ \mathrm{Im}(\vec{\Gamma}) = \mathrm{Im}(\vec{\Gamma})$$

and for $E \in \Sigma$, $\mathrm{Im}(\vec{\Gamma})$ is not the zero vector. Hence, $|\Gamma|^2 S$ has an eigenvalue 1 which proves (A.1). □

Appendix B. Some identities for Gaussian integrals and the Fourier transform

The following identities are used at various parts in the article.

Lemma B.1. *Let D be an invertible, symmetric $k \times k$ matrix with positive definite real part, i.e. $D = D^\top$, $\mathrm{Re}(D) > 0$. Then, for any complex vector $v \in \mathbb{C}^k$ one has the Gaussian integral*

$$\int_{\mathbb{R}^k} e^{-\frac{1}{2}(x+v)\cdot D(x+v)}\, d^k x \;=\; \frac{(2\pi)^{k/2}}{\sqrt{\det(D)}}\,. \tag{B.1}$$

Some care needs to be taken to select the correct branch of $\sqrt{\det(D)}$. If $D = A + iB$ where $A > 0$ is the real part, then we write $D = \sqrt{A}(1 + iA^{-1/2}BA^{-1/2})\sqrt{A}$ where $\sqrt{A}$ has the same eigenspaces as A and the corresponding eigenvalues are the positive square roots of the eigenvalues of A. Furthermore, $A^{-1/2}BA^{-1/2}$ is diagonalizable by a real orthogonal matrix. This diagonalizes $1 + iA^{-1/2}BA^{-1/2}$ as well and the eigenvalues have all real part 1. Hence, we may define $\sqrt{1 + iA^{-1/2}BA^{-1/2}}$ by taking the same eigenspaces and the principal branch of the square roots of the eigenvalues. Then (B.1) *is correct with $\sqrt{\det(D)} = \det(\sqrt{A})\det(\sqrt{1 + iA^{-1/2}BA^{-1/2}})$.*

Proof. In one dimension one has the well known integral formula

$$\int_{-\infty}^{\infty} e^{-z(x+c)^2}\, dx = \frac{\sqrt{\pi}}{\sqrt{z}} \tag{B.2}$$

for $\mathrm{Re}(z) > 0$, where the square root is the principal branch and c is any fixed complex number. Now if $D = A + iB$, then use a basis change $y = O\sqrt{A}x$, where O is a real orthogonal matrix such that $OA^{-1/2}BA^{-1/2}O^\top$ is diagonal. This leads to a Gaussian integral with a diagonal matrix and then (B.1) follows from (B.2). $\square$

Recall that $\mathcal{M}(g(x))$ denoted the multiplication operator that multiplies by $g(x)$. Similarly, let $\mathcal{C}(g(x))$ denote the convolution operator, i.e.

$$\mathcal{C}(g(x))f\,(y) = \int g(y-x)f(x)\, d^k x$$

Lemma B.2. *Let $\mathcal{F}$ denote the Fourier transform on $\mathbb{R}^k$, and let D be a symmetric, invertible $k \times k$ matrix with positive semi-definite imaginary part $\mathrm{Im}(D) \geq 0$. Then as a map from $L^1(\mathbb{R}^k) \cap L^1(\mathbb{R}^k)$ to $L^2(\mathbb{R}^k)$ one has*

$$\mathcal{F}^* \mathcal{M}(e^{\frac{1}{2}ix\cdot Dx})\mathcal{F} \;=\; \frac{\mathcal{C}(e^{-\frac{1}{2}x\cdot iD^{-1}x})}{(2\pi)^{k/2}\sqrt{\det(-iD)}} \tag{B.3}$$

where $\sqrt{\det(-iD)}$ is selected as in Lemma B.1 (note that $\mathrm{Re}(-iD) \geq 0$).

If D is real, i.e. $\mathrm{Im}(D) = 0$, this can be re-written as

$$\mathcal{F}^* \mathcal{M}(e^{\frac{1}{2}ix\cdot Dx})\mathcal{F} \;=\; \frac{1}{\sqrt{\det(-iD)}}\, \mathcal{M}(e^{-\frac{1}{2}ix\cdot D^{-1}x})\, \mathcal{S}_D\, \mathcal{F}\, \mathcal{M}(e^{-\frac{1}{2}ix\cdot D^{-1}x}) \tag{B.4}$$

$\mathcal{S}_D$ denotes the group action $\mathcal{S}_D f(y) = f(D^{-1}y)$.

Proof. First assume $\mathrm{Im}(D) > 0$, then for $f \in L^1(\mathbb{R}^k)$ and any $y \in \mathbb{R}^k$, the map $(x,w) \mapsto e^{-iy\cdot x} e^{\frac{1}{2}ix\cdot Dx} e^{ix\cdot w} f(w)$ is in $L^1(\mathbb{R}^{2k})$ and one finds

$$\begin{aligned}(2\pi)^k \mathcal{F}^* \mathcal{M}(e^{\frac{1}{2}ix\cdot Dx})\mathcal{F}f)(y) &= \int e^{-iy\cdot x} e^{\frac{1}{2}ix\cdot Dx} e^{ix\cdot w} f(w)\, d^k w\; d^k x \\ &= \int \left[\int e^{\frac{1}{2}[x+D^{-1}(w-y)]\,\cdot\, iD[x+D^{-1}(w-y)]}\, d^k x \right] \, e^{-\frac{i}{2}(y-w)\cdot D^{-1}(y-w)}\, f(w)\, d^k w \\ &= \frac{(2\pi)^{k/2}}{\sqrt{\det(-iD)}}\; \mathcal{C}(e^{-\frac{i}{2}x\cdot D^{-1}x})\, f\,(y)\,. \end{aligned} \tag{B.5}$$

Now, if $\mathrm{Im}(D)$ is only positive semi-definite, we approach D by $D + i\epsilon$ and let $f \in L^1(\mathbb{R}^k) \cap L^2(\mathbb{R}^k)$. Then as $\epsilon \downarrow 0$, the right hand side converges point wise (for fixed y). As the L^2 norm is uniformly bounded by $\|f\|_2$, Dominated Convergence shows convergence in $L^2(\mathbb{R}^k)$. As the operators $\mathcal{M}(e^{\frac{1}{2}ix\cdot(D+i\epsilon)x})$ converge for $\epsilon \downarrow 0$ in the strong operator topology, we also get convergence on the left hand side in $L^2(\mathbb{R}^k)$.

For (B.4) note

$$\int e^{-\frac{i}{2}(y-w)\cdot D^{-1}(y-w)}\, f(w)\, d^k w \;=\; e^{-\frac{i}{2}y\cdot D^{-1}y} \int e^{iD^{-1}y\cdot w} e^{-\frac{i}{2}w\cdot{}^{-1}w} f(w) d^k w\;.$$

□

Equation (B.3) yields the following.

Corollary B.3. *Let $\mathbb{S}_+$ be the the set of invertible, symmetric $k \times k$ matrices with positive semi-definite imaginary part. The map $F(D,f) = \mathcal{F}^*\mathcal{M}(e^{\frac{1}{2}ix\cdot Dx})\mathcal{F}f$ maps $\mathbb{S}_+ \times (L^1(\mathbb{R}^k) \cap L^2(\mathbb{R}^k)$ to $L^2(\mathbb{R}^k) \cap L^\infty(\mathbb{R}^k)$, and satisfies*

$$\|F(D,f)\|_2 \le \|f\|_2\,, \quad \|F(D,f)\|_\infty \le \frac{\|f\|_1}{(2\pi)^{k/2}\sqrt{|\det(D)|}}\,. \tag{B.6}$$

Moreover, if $D_n \to D$ in $\mathbb{S}_+$ and $f_n \to f$ in $L^1(\mathbb{R}^k)$ then by Dominated Convergence, $F(D_n, f_n)$ converges point wise, i.e. $\lim_{n\to\infty} F(D_n,f_n)(y) = F(D,f)(y)$.

References

[Aiz] M. Aizenman, *Localization at weak disorder: some elementary bounds*, Rev. Math. Phys. **6**, 1163-1182 (1994)

[ASW] M. Aizenman, R. Sims and S. Warzel, *Stability of the absolutely continuous spectrum of random Schrödinger operators on tree graphs*, Prob. Theor. Rel. Fields, **136**, 363-394 (2006)

[AM] M. Aizenman and S. Molchanov, *Localization at large disorder and extreme energies: an elementary derivation*, Commun. Math. Phys. **157**, 245-278 (1993)

[AW] M. Aizenman and S. Warzel, *Resonant delocalization for random Schrdinger operators on tree graphs*, preprint arXiv:1104.0969 (2011)

[AW2] M. Aizenman and S. Warzel, *Absolutely continuous spectrum implies ballistic transport for quantum particles in a random potential on tree graphs*, J. Math. Phys. **53**, 095205 (2012)

[Ber] F.A. Berezin, *The method of second quantization*, Academic Press, New York, 1966

[Breu] J. Breuer, *Localization for the Anderson model on trees with finite dimensions*, Ann. Henri Poincarè **8**, 1507-1520 (2007)

[CKM] R. Carmona, A. Klein and F. Martinelli, *Anderson localization for Bernoulli and other singular potentials*, Commun. Math. Phys. **108**, 41-66 (1987)

[DLS] F. Delyon, Y. Levy and B. Souillard, *Anderson localization for multidimensional systems at large disorder or low energy*, Commun. Math. Phys. **100**, 463-470 (1985)

[DK] H. von Dreifus and A. Klein, *A new proof of localization in the Anderson tight binding model*, Commun. Math. Phys. **124**, 285-299 (1989)

[Efe] K. B. Efetov, *Supersymmetry and theory of disordered metals*, Advances Phys. **32**, 53-127 (1983)

[FHH] R. Froese, F. Halasan and D. Hasler, *Absolutely continuous spectrum for the Anderson model on a product of a tree with a finite graph*, arXiv:1008.2949v1 (2010)

[FHS] R. Froese, D. Hasler and W. Spitzer, *Absolutely continuous spectrum for the Anderson Model on a tree: A geometric proof of Klein's Theorem*, Commun. Math. Phys., **269**, 239-257 (2007)

[FHS2] R. Froese, D. Hasler and W. Spitzer, *Absolutely continuous spectrum for a random potential on a tree with strong transverse correlations and large weighted loops*, Rev. Math. Phys. **21**, 709-733 (2009)

[FMSS] J. Fröhlich, F. Martinelli, E. Scoppola and T. Spencer *Constructive proof of localization in the Anderson tight binding model*, Commun. Math. Phys. **101**, 21-46 (1985)

[FS] J. Fröhlich and T. Spencer, *Absence of diffusion in the Anderson tight binding model for large disorder or low energy*, Commun. Math. Phys. **88**, 151-184 (1983)

[GMP] Ya. Gol'dsheid, S. Molchanov and L. Pastur, *Pure point spectrum of stochastic one dimensional Schrödinger operators*, Funct. Anal. Appl. **11**, 1-10 (1977)

[Hal] F. Halasan, *Absolutely continuous spectrum for the Anderson model on trees*, PhD thesis 2009, arXiv:0810.2516v3 (2008)

[Ka] T. Kato, *Wave operators and similarity for some non self-adjoint operators*, Mat. Ann. **162**, 258-279 (1966)

[Kel] M. Keller, *On the spectral theory of operators on trees*, PhD Thesis 2010, accessible at arXiv:1101.2975

[KLW] M. Keller, D. Lenz and S. Warzel, *On the spectral theory of trees with finite cone type*, arXiv:1001.3600v2 (2011), to appear in Israel J. Math.

[KLW2] M. Keller, D. Lenz and S. Warzel, *Absolutely continuous spectrum for random operators on trees of finite cone type*, arXiv:1108.0057 (2011)

[Kl1] A. Klein, *The supersymmetric replica trick and smoothness of the density of states for random Schrodinger operators*, Proc. Symposia in Pure Mathematics **51**, 315-331 (1990)

[Kl2] A. Klein, *Localization in the Anderson model with long range hopping*, Braz. J. Phys. **23**, 363-371 (1993)

[Kl3] A. Klein, *Absolutely continuous spectrum in the Anderson model on the Bethe lattice*, Math. Res. Lett. **1**, 399-407 (1994)

[Kl4] A. Klein, *Absolutely continuous spectrum in random Schrödinger operators*, Quantization, nonlinear partial differential equations, and operator algebra (Cambridge, MA, 1994), 139-147, Proc. Sympos. Pure Math. **59**, Amer. Math. Soc., Providence, RI, 1996

[Kl5] A. Klein, *Spreading of wave packets in the Anderson model on the Bethe lattice*, Commun. Math. Phys. **177**, 755–773 (1996)

[Kl6] A. Klein, *Extended states in the Anderson model on the Bethe lattice*, Advances in Math. **133**, 163-184 (1998)

[KLS] A. Klein, J. Lacroix and A. Speis, *Localization for the Anderson model on a strip with singular potentials*, J. Funct. Anal. **94**, 135-155 (1990)

[KS] A. Klein and C. Sadel, *Absolutely Continuous Spectrum for Random Schrödinger Operators on the Bethe Strip*, Math. Nachr. **285**, 5-26 (2012)

[KS2] A. Klein and C. Sadel, *Ballistic Behavior for Random Schrödinger Operators on the Bethe Strip*, J. Spectr. Theory **1**, 409-442 (2011)

[KSp] A. Klein and A. Speis, *Smoothness of the density of states in the Anderson model on a one-dimensional strip*, Annals of Phys. **183**, 352-398 (1988)

[Klo] F. Klopp, *Weak disorder localization and Lifshitz tails*, Commun. Math. Phys. **232**, 125-155 (2002)

[KuS] H. Kunz and B. Souillard, *Sur le spectre des operateurs aux differences finies aleatoires*, Commun. Math. Phys. **78**, 201-246 (1980)

[Lac] J. Lacroix, *Localisation pour l'opérateur de Schrödinger aléatoire dans un ruban*, Ann. Inst. H. Poincaré ser **A40**, 97-116 (1984)

[Nir] L. Nirenberg, *Topics in Nonlinear Functional Analysis*, New York: Courant Institute of Mathematical Sciences, 1974

[Sad] C. Sadel, *Absolutely continuous spectrum for random Schrödinger operators on tree-strips of finite cone type*, Annales Henri Poincare, **14**, 737-773 (2013)

[SS] C. Sadel and H. Schulz-Baldes, *Random Dirac Operators with time reversal symmetry*, Commun. Math. Phys. **295**, 209-242 (2010)

[SW] B. Simon and T. Wolff, *Singular continuum spectrum under rank one perturbations and localization for random Hamiltonians*, Commun. Pure. Appl. Math. **39**, 75-90 (1986)

[Wa] W.-M Wang, *Localization and universality of Poisson statistics for the multidimensional Anderson model at weak disorder*, Invent. Math. **146**, 365-398 (2001)

Mathematics Department, University of British Columbia, Vancouver, BC, V6T 1Z2, Canada

E-mail address: `csadel@math.ubc.ca`